



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

**0 342 682
A2**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 89108976.5

(51) Int. Cl.⁴: **C07D 209/28 , C07C 83/10 ,
A61K 31/405 , A61K 31/16**

(22) Date of filing: 18.05.89

Claims for the following Contracting States: ES
+ GR.

(30) Priority: 19.05.88 US 196584

(43) Date of publication of application:
23.11.89 Bulletin 89/47

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(71) Applicant: **WARNER-LAMBERT COMPANY**
201 Tabor Road
Morris Plains New Jersey 07950(US)

(72) Inventor: Cetenko, Wlaczslaw Antin
3565 Green Brier
Ann Arbor Michigan 48105(US)
Inventor: Connor, David Thomas
2453 Antietam
Ann Arbor Michigan 48105(US)
Inventor: Flynn, Daniel Lee
17 North Bristol Court
Mundelein Illinois 60060(US)
Inventor: Sircar, Jagadish Chandra
3615 Charter Place
Ann Arbor, Michigan 48105(US)

(74) Representative: Mansmann, Ivo
c/o Gödecke AG - Patentabteilung Postfach
569 - Mooswaldallee 1-9
D-7800 Freiburg(DE)

(54) **Novel hydroxamate derivatives of selected nonsteroidal anti-inflammatory acyl residues having cyclooxygenase and 5-lipoxygenase inhibition.**

(57) The present invention is novel selected hydroxamic acid derivatives of acyl residues of selected NSAIDS, i.e. having 5-lipoxygenase and cyclooxygenase inhibiting properties, pharmaceutical compositions for treating conditions advantageously affected by the inhibition and methods for treating these conditions in mammals, including humans suffering therefor.

EP 0 342 682 A2


NOVEL HYDROXAMATE DERIVATIVES OF SELECTED NONSTEROIDAL ANTIINFLAMMATORY ACYL RESIDUES HAVING CYCLOOXYGENASE AND 5-LIPOXYGENASE INHIBITION

BACKGROUND OF THE INVENTION

The present invention is novel hydroxamate analogs of acyl residues of selected nonsteroidal antiinflammatory drug(s) (NSAID(S)). Such NSAIDS include indomethacin and its analogs of U.S. Patent No. 3,161,654; sulindac and its analogs of U.S. Patent No. 3,654,349; tolmetin and its analogs of U.S. Patent No. 3,752,826; and furofenac and its analogs of U.S. Patent No. 4,029,811; fentiazac and its analogs of U.S. Patent No. 3,476,766; clidanac and its analogs of U.S. Patent No. 3,565,943; ketorolac of U.S. Patent No. 4,089,969; oxepinac of British Patent No. 1,476,214; fenclorac of U.S. Patent No. 3,864,384; lonazolac of U.S. Patent No. 4,146,721; metiazinic acid of U.S. Patent Nos. 3,455,917 and 3,424,748; clopirac of Belgian Patent No. 777,207; clometacine of British Patent No. 1,260,868; etodolac of U.S. Patent No. 3,939,178; indoprofen of British Patent No. 1,344,663; piroprofen of U.S. Patent No. 3,641,040; carprofen of U.S. Patent No. 3,896,145; oxaprozin of U.S. Patent No. 3,578,671; pranoprofen of U.S. Patent No. 3,931,205; suprofen of U.S. Patent No. 4,035,376; miroprofen of U.S. Patent No. 3,978,071; tiroxaprofen of U.S. Patent No. 3,933,840; furaprofen of German Patent No. DE 3026402 or furobufen of U.S. Patent No. 3,728,349 or diclofenac of U.S. Patent No. 3,558,690; and bucloxic acid of U.S. Patent No. 3,754,021.

The above patents are each incorporated by reference.


Anthranilic acid derivatives of a fenamic acid series are previously known to include the generic compounds of the British Patent No. 989,951 or formula



wherein Ar is α,α,α -trifluoro-m-tolyl; 2,3-xylyl, or 2,6-dichloro-m-tolyl, also in U.S. Patent No. 3,852,333.

Of these, disclosures for unsubstituted hydroxamic acid of the acetyl residue of indomethacin are found in U.S. Patent No. 3,624,103 and diclofenac are found in U.S. Patent Nos. 4,092,430 and 4,173,577. However, no teaching to further substituted hydroxamic acids of indomethacin and diclofenac having the activity of the present invention compounds is found.

Among related aminobenzhydroxamic acids also previously known are compounds of the formula




wherein R' is a saturated fatty hydrocarbon radical; phenyl, phenylalkyl, wherein the rings are optionally substituted by lower alkyl or lower alkoxy; or an aromatic heterocyclic group. This disclosure is in Japanese Application 24578/67 filed April 2, 1964 by the Takeda Chemical Industry Co., Ltd. as an o-aminobenzhydroxamic acid analgesic derivative having less toxicity and analgesic, antitarrhic, and antifebrile activity.

Other related disclosures include U.S. Patent No. 4,029,815 to compounds of the formula

5

10




wherein X_5 is trifluoromethyl, difluoromethyl, or nitro, X_4 is H, Br, Cl, or nitro, and Q may be NHOH. These compounds have utility as antidiarrheal agents.

Selected fenamic acid derivatives having an hydroxamic acid derived substituent are found in copending U.S. application Serial Number 134725.

Cyclized α -aminobenzhydroxamic-O-methylether of the formula

20

25




wherein R'' is alkyl, aralkyl, or a basic side chain and R' is H, Cl, or Br; are disclosed by Wolf, E. and Kohl, H. in "Cyclisierungsreaktionen von am Aminostickstoff Substituierten α -Aminobenzhydroxamsaure-O-methylesteren," *Ann. Chem. Liebigs*, 1975, 1245-1251.

Wolf and Kohl also disclose an intermediate hydroxamic acid derivative from which the cyclized α -aminobenzhydroxamic-O-methylether are made. The intermediate is

35

40




wherein R_c , R_a , and R_d are as defined below.


Other cyclized α -aminobenzhydroxamic acids disclosed are

45

50




or



wherein R_a is $\text{CH}_2\text{CO}_2\text{C}_2\text{H}_5$, $\text{C}_6\text{H}_3\text{Cl(p)NO}_2(\text{m})$ and suggesting that R_a may also be $\text{SO}_2\text{C}_6\text{H}_4\text{CH}_3(\text{p})$; and

55




wherein R_b is H or phenyl; R_c is H, $\text{CH}_2\text{C}_6\text{H}_5$, $\text{C}_6\text{H}_4\text{Cl(p)}$, $\text{CH}_2\text{C}_6\text{H}_4\text{Cl(p)}$, C_6H_5 , or CH_3 ; R_e is H or NO_2 ; and R_d is H or Cl. However, Wolf and Kohl do not disclose activity for these cyclized compounds and, further, do not make obvious the present invention.

Broadly, hydroxamic acid derivatives of selected aryl ring systems are disclosed in European Application Publication No. 0 196 184 and 196 674 having surprisingly high potency particularly by inhalation, oral efficacy, and with a surprisingly long duration of action. However, these aryl ring systems are in no way related to the present NSAID type compounds.

Two disclosures by Summers et al, (1) *J. Med. Chem.*, 1987, 30, 574-80 and 2121-2126 and (2) *In Vivo Characterization of Hydroxamic Acid Inhibitors of 5-Lipoxygenase* disclosed at a Poster session at a National ACS meeting (New Orleans) meeting in September, 1987 (Abstract) disclose hydroxamic acids as inhibitors of 5-lipoxygenase, however, the disclosures do not extend beyond very limited representative examples not including any derivatives having the selected acyl residues of the present invention.

Thus, the present invention are to selected novel derivatives of NSAIDS and pharmaceutically acceptable acid addition or base salts thereof, pharmaceutical compositions for treating allergy, psoriasis, inflammation, arthritis, pain, pyrrhia, and the methods for such treatment.


The present invention is a novel compound of the formula (I)




I

and pharmaceutically acceptable acid addition and base salts thereof; wherein

- (i) $\text{R}-\overset{\text{O}}{\parallel}{\text{C}}$ is an acyl residue of selected NSAIDS;
- (ii) R_1 is hydrogen, lower alkyl, or acyl; and
- (iii) R_2 is H, lower alkyl, cycloalkyl of from three to twelve carbon atoms of which from three to seven are ring carbons, aryl, aralkyl or heteroaryl with the overall proviso that when the acyl residue is of the formula



wherein R_4 is hydrogen or lower alkyl, then R_3 is hydrogen, phenylmethyl, allyl, vinyl, isopropenyl, benzoyl, chlorobenzoyl, methoxybenzoyl, and thiomethylbenzoyl; and when the acyl residue is of the formula



wherein R_5 , R_6 , R_7 and R_8 are independently H, chlorine, fluorine, or bromine, lower alkyl, lower alkoxy, or trifluoromethyl but when R_4 is hydrogen then are not all hydrogen; and R_4 is otherwise independently as defined above; then R_1 and R_2 cannot both be hydrogen.

The present invention is also a pharmaceutical composition for the treatment of conditions advantageously affected by the inhibition of 5-lipoxygenase and/or cyclooxygenase which comprises administering an amount effective for inhibiting 5-lipoxygenase and/or cyclooxygenase of a novel compound of the formula I as defined above; and a pharmaceutically acceptable carrier.


Further, the present invention also provides a method of use for a composition of a compound of the formula I, as defined hereinbefore, or physiologically acceptable acid addition or base salt thereof for the manufacturing of pharmaceuticals for the treatment of a condition which is advantageously affected by the inhibition of the lipoxygenase and cyclooxygenase enzymes of the mammalian, including human, arachidonic acid metabolism, which method comprises inhibition of such enzymes by administration to a mammal of a lipoxygenase and cyclooxygenase inhibiting amount of any such compound or salt in unit dosage form.

Further, the present invention also provides any compound or composition of the formula I' or physiologically acceptable salt thereof, for the manufacturing of pharmaceuticals, for use as a medical therapeutic or prophylactic agent, to methods of medical therapeutic or prophylactic treatment by administration to a mammal of a medically therapeutic or prophylactic effective amount of any such compound or salt. The kinds of medical therapy and prophylaxis pertinent to the foregoing and therefore in that sense comprising part of the present invention, are elaborated by way of example in the following paragraphs which are not intended to be construed as in any way limiting the scope of these aspects of said invention.

In the compounds of formula I the term "lower alkyl" is meant to include a straight or branched alkyl group having one to six carbon atoms, such as, for example, methyl, ethyl, propyl, butyl, pentyl or hexyl and isomers thereof.

Lower alkoxy is O-alkyl or of from one to four carbon atoms as defined above for "lower alkyl".

Acyl is a



45 $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}-\text{phenyl optionally substituted by a lower alkyl, fluoro, chloro, bromo, trifluoromethyl, hydroxy, or lower alkoxy; and wherein the lower alkyl is as defined above.} \end{array}$

Q Cycloalkyl of from three to ten carbons having from three to seven ring carbons includes cyclopropyl, cyclobutyl, methylcyclobutyl, ethylcyclobutyl, dimethylcyclobutyl, cyclopentyl, and the like.

Aryl or aromatic radical is phenyl unsubstituted and substituted with from one to three substituents selected from the group consisting of hydroxy, lower alkoxy, fluoro, chloro, bromo, trifluoromethyl, lower alkyl, CN, $-\text{S}(\text{O})_n$ -lower alkyl wherein n is zero to four NO_2 , or NR_9R_{10} wherein R_9 and R_{10} are independently hydrogen or lower alkyl.

55 An aralkyl is an aryl as defined above attached through a lower alkylene where the alkylene, straight or branched chain, is of from one to four carbons such as methylene, 1,2-ethylene, 1,1-ethylene, propylene, and the like.


Heterocycles includes derivatives of furan, thiophene, pyrrole, piperidine, dihydrofuran, pyridine, thiazole, piperazine, oxazole, benzofuran, tetrahydroquinoline, quinoline, indole, dihydroindole, benzothiofene, dihydrobenzothiofene, benzoxazole, and similar heterocyclic rings.

The cycloalkyl and fused cycloalkyl groups may be mono or polycyclic and contain from three to twenty carbons. These groups include cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, bornyl, norbornyl, indanyl and the like which are optionally substituted by lower alkyl up to a total of twelve carbons.

Physiologically acceptable salts are meant to be synonymous with pharmacologically acceptable salts in this invention.

An acyl residue of selected NSAIDS is RCO of the formula

(a)



I_a

in which

R_{1a} is selected from the group consisting of benzene, naphthalene, biphenyl and substituted benzene, naphthalene and biphenyl radicals in which said substituent is selected from the group consisting of halogen, lower alkyl, lower alkylthio, lower alkoxy, trifluoromethyl, phenoxy, lower alkyl phenoxy, lower alkoxy phenoxy, halogenophenoxy, trifluoroacetyl, difluoroacetyl, monofluoroacetyl, di-lower alkyl sulfamyl, lower alkanoyl, di-lower alkyl carboxamido, cyano, carb-lower alkoxy, trifluoromethylthio, lower alkyl sulfinyl, lower alkylsulfonyl, benzylthio, lower alkylbenzylthio, lower alkoxybenzylthio, halogenobenzylthio, mercapto, nitro, amino, di-(lower alkyl)amino, lower alkylamino, lower alkanoylamino, hydroxy, lower alkanoyloxy, trifluoroacetoxy, difluoroacetoxy, monofluoroacetoxy, benzyloxy, lower alkylbenzyloxy, lower alkoxybenzyloxy, and halogenobenzoyloxy;

R_{2a} is selected from the group consisting of hydrogen, lower alkenyl and lower alkyl;


R_{3a} is selected from the group consisting of hydrogen and lower alkyl;

R_{4a} is selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, fluorine and trifluoromethyl;

R_{5a} is selected from the group consisting of hydrogen, hydroxy, lower alkyl, lower alkoxy, nitro, amino, lower alkylamino, di(lower alkyl) amino, lower alkanoylamino, lower alkanoyl, lower alkylamino, bis(hydroxy lower alkyl)amino, 1-pyrrolidino, 4-methyl-1-piperiziny, 4-morpholinyl, cyano, amino lower alkyl, di-lower alkyl amino, lower alkyl, trifluoromethyl, halogen, di(lower alkyl)sulfamyl, benzylthio, lower alkylbenzylthio, lower alkoxybenzylthio, halogenobenzylthio, benzyloxy, lower alkylbenzyloxy, lower alkoxybenzyloxy, halogenobenzoyloxy, lower alkenyl, lower alkenyloxy, 1-azacyclopropyl, cyclopropyl(lower alkoxy) methoxy, and cyclobutyl(lower alkoxy)methoxy.

(The most preferred compound of the present invention is the compound of formula I_a wherein RCO is the acyl residue of 1-(4-chlorobenzoyl)-2-methyl-5-methoxy-3-indole acetic acid.)

(b)



wherein:

R_{1b} is hydrogen, lower alkyl or halogenated lower alkyl;

20 R_{2b} is hydrogen or alkyl;

R_{3b}, R_{4b}, R_{5b} and R_{6b} each are hydrogen, lower alkyl, lower alkoxy, nitro, amino, lower alkylamino, lower dialkylamino, lower dialkylamino lower alkyl, sulfamyl, lower alkylthio, mercapto, hydroxy, hydroxy lower alkyl, lower alkylsulfonyl, halogen, carboxyl, carbo-lower alkoxy, carbamido, halogenoalkyl, cycloalkyl, or cycloalkoxy;

25 R_{7b} is lower alkylthio, alkylsulfinyl or alkylsulfonyl;

R_{8b} is hydrogen, halogen, hydroxy, alkoxy or haloalkyl.

(The most preferred compound of the formula I wherein RCO is defined in group (b) is the compound wherein RCO is I_b wherein R_{1b} is hydrogen; R_{2b} is methyl; R_{3b} is hydrogen; R_{4b} is fluoro; R_{5b} is hydrogen; R_{6b} is hydrogen; R_{7b} is p-methylsulfinyl; and R_{8b} is hydrogen.)

30

35



40

45


50

55

(c)



25 and



40 wherein:

Ar_c represents a member selected from the group consisting of phenyl, thienyl, 5-methylthienyl, monosubstituted phenyl, disubstituted phenyl and trisubstituted phenyl, each substituent of said substituted phenyls being a member selected from the group consisting of halo, lower alkyl, trifluoromethyl, lower alkoxy, nitro, amino, methylthio and cyano;

45 Ar_{1c} represents a member selected from the group consisting of phenyl, monosubstituted phenyl, disubstituted phenyl and trisubstituted phenyl, each substituent of said substituted phenyls being a member selected from the group consisting of halo, lower alkyl and lower alkoxy;

R_c represents a member selected from the group consisting of hydrogen and lower alkyl;

R_{1c} represents a member selected from the group consisting of hydrogen, lower alkyl and benzyl;

50 R_{2c} represents $\text{C}(=\text{O})-$;

R_{3c} represents $\text{C}(=\text{O})-$;


R_{4c} represents lower alkyl;

55 R_{5c} represents lower alkyl; and

R_{6c} represents a member selected from the group consisting of hydrogen and lower alkyl.


(The most preferred compound of the formula I wherein RCO is defined in group (c) is the compound wherein RCO is the acyl residue of 5-(4-methylbenzoyl)-1-methylpyrrolo-2-acetic acid.)

(d)




wherein R_d represents a group selected from:

5



and


10



15

(e)

20

 I_e

25


wherein R^{1e} and R^{2e} are each radicals of the group consisting of thienyl, furyl, naphthyl, phenyl, and phenyl bearing from one to two substituents of the group consisting of (lower)alkyl, (lower)alkoxy, chloro, bromo, fluoro, di-(lower alkyl)amino, nitro, amino and trifluoromethyl, and wherein R^{3e} is a radical from the group consisting of $-CH_2-CO-$, $-CH_2-CH_2-CO-$, and $-CH(CH_3)-CO-$.

(The most preferred compound of the formula I wherein RCO is defined in group (e) is the compound wherein RCO is I_e wherein R^{1e} is phenyl, R^{3e} is $-CH_2CO-$ and R^{2e} is p-chlorophenyl.)

35

(f)

40

 I_f

45


wherein Y is hydrogen, chloro, bromo, fluoro, hydroxy, (lower)alkyl, (lower)alkoxy, mercapto, cyano, nitro, amino or (lower)alkylthio.

(The most preferred compound of the formula I wherein RCO is defined in group (f) is the compound wherein RCO is the acyl residue of (\pm)-6-chloro-5-cyclohexyl-1-indancarboxylic acid.)

50

55


(g)



wherein R_g represents hydrogen or a lower alkyl group having from one to four carbon atoms and R_{1g} represents hydrogen, a lower alkyl group having from one to 4 carbon atoms, a lower alkoxy group having from one to four carbon atoms, chloro, fluoro or bromo, the R_{1g} substitution being at the ortho, meta or para positions of the aroyl group.

(The most preferred compound of the formula I wherein RCO is defined in group (g) is the compound wherein RCO is the acyl residue of 5-benzoyl-1,2-dihydro-3H-pyrrolo-[1,2-a]pyrrole-1-carboxylic acid.)


(h)



30 wherein

R_{1h} represents a hydrogen or halogen atom or a trihalomethyl or lower alkoxy group; R_{2h} represents a hydrogen atom or a lower alkyl group;

(i)



45 wherein

R_{1i} is a member selected from the group consisting of cycloalkyl with five to seven carbon atoms and lower alkyl substituted cycloalkyl;

R_{2i} is a member selected from the group consisting of halogen, nitro, cyano, trifluoromethyl and lower alkylsulfonyl;


R_{3i} is hydrogen;

X_1 is halogen.

(The most preferred compound of the formula I wherein RCO is defined in group (i) is the compound wherein RCO is the acyl residue of α ,m-dichloro-p-cyclohexylphenylacetic acid.)

(j)


5

I_{ji}

10

wherein R^{1ji} and R^{2ji} are selected from the group consisting of phenyl, halophenyl, lower alkylphenyl, dimethylphenyl, lower alkoxyphenyl, dimethoxyphenyl, lower alkylmercaptophenyl, trifluoromethylphenyl, furyl, thienyl, and naphthyl with the proviso that at least one of R^{1ji} and R^{2ji} is phenyl or substituted phenyl;

15


I_{jii}

20

wherein R^{1jii} is hydrogen, lower alkyl, phenyl, phenyl monosubstituted with lower alkyl, lower alkoxy or halogen, naphthyl, benzyl or benzyl wherein the phenyl ring thereof is monosubstituted with lower alkoxy or halogen;

25


30

I_{jiii}

35

wherein R^{1jiii} is phenyl; R^{2jiii} is phenyl, p-halogenophenyl, p-methylphenyl or furyl; R^{3jiii} is hydrogen, phenyl, or furyl; or

40

I_{jiv}

45


wherein R^{1jiv} is phenyl, R^{2jiv} is phenyl, p-halogenophenyl, p-methylphenyl, R^{3jiv} is hydrogen or phenyl.

(The most preferred compound of the formula I wherein RCO is defined in group (j) is the compound wherein RCO is the acyl residue of 1-phenyl-3-(p-chlorophenyl)pyrazol-4-yl-acetic acid.)

50

55


(k)

 I_k

wherein R_k represents hydrogen or methyl, R_{1k} represents hydrogen, methyl or ethyl, and Y_k represents hydrogen, halogen, or alkyl or alkoxy of 1 through 4 carbon atoms.

(The preferred compounds of the formula I wherein RCO is defined in group (k) is the compound having I_k wherein R_k is hydrogen or methyl, R_{1k} is hydrogen, methyl or ethyl and Y_k is hydrogen or methoxy, and the most preferred is the compound having I_k wherein R_k is methyl, R_{1k} is hydrogen and Y_k is hydrogen.)


(1)

 I_1

wherein R_{11} is a lower alkyl or lower alkyl substituted by cycloalkyl of from three to 12 carbons having from three to seven ring carbons, lower alkenylenyl, lower alkynylenyl, aryl, benzoyl, benzoyl substituted by halogen, lower alkyl, lower alkoxy or heterocycle, such as morpholinyl piperidinyl, furanyl, and thiophenyl; R_{2a} and R_{3a} are the same or different and are lower alkyl, aryl and R_{4a} is hydrogen or lower alkyl.

(The most preferred compound of the formula I wherein RCO is defined in group (1) is the compound wherein RCO is the group I_1 wherein R_{21} and R_{31} are CH_3 , R_{41} is hydrogen and R_1 is p-chlorophenyl.)

(m)

 I_m

[in which A_m represents a lower straight chain or branched alkylene group;

R_m represents a cyclohexyl or aromatic radical;


R'_m represents a halogen atom, trifluoromethyl radical, an alkoxy or alkyl radical containing 1 to 4 carbon atoms or a N,N-dialkylamino radical wherein each of the alkyl groups contain 1 to 4 carbon atoms] as well as alkyl esters thereof (in which the alkyl group in the alcoholic moiety contains 1 to 4 carbon atoms).

(The most preferred compound of the formula I wherein RCO is defined in group (m) is the compound wherein RCO is the group I_m wherein R'_m is methoxy, A_m is $-CH_2-$, and R_m is p-chlorophenyl.)

(n)


5

10

 I_n

in which R^{1n} is selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower cycloalkyl, phenyl, benzyl and 2-thienyl, R^{2n} , R^{3n} , R^{4n} and R^{5n} are the same or different and are each selected from the group consisting of hydrogen and lower alkyl, R^{6an} , R^{6bn} , R^{6cn} , and R^{6dn} are the same or different and selected from the group consisting of hydrogen, lower alkyl, hydroxy, lower alkoxy, benzyloxy, lower alkanoyloxy, nitro, halo, mercapto, lower alkylthio, trifluoromethyl, amino and sulfamoyl, R^{7n} is selected from the group consisting of hydrogen, lower alkyl and lower alkenyl, X is selected from the group consisting of oxy and thio, Y is selected from the group consisting of carbonyl,

25




which each of R^{8n} , R^{9n} , R^{10n} , R^{11n} , R^{12n} and R^{13n} is hydrogen or lower alkyl.

(The most preferred compound of the formula I wherein RCO is defined in group (n) is the compound wherein RCO is the acyl residue of 1,8-diethyl-1,3,4,9-tetrahydropyrano[3,4-b]indole-1-acetic acid.)

(o)

35

40

 I_o


wherein R_o is hydrogen or lower alkyl.

(The preferred compounds of the formula I wherein RCO is defined in group (o) is the compound I_o wherein R_o is CH_3 .)

45

(p)

50

 I_p


in which R_{1p} is hydrogen or lower alkyl; R_{2p} is hydrogen, alkyl, alkenyl three or four ring-membered cycloalkyl, cycloalkenyl or cycloalkyl-methyl cycloalkenyl-methyl; Ph is (i) 1,3- or 1,4-phenylene, (ii) (lower alkyl)-1,3- or 1,4-phenylene, (iii) (lower alkoxy)-1,3- or 1,4-phenylene, (iv) mono- or di-(halogeno)-1,3- or 1,4-phenylene, (v) (trifluoromethyl)-1,3- or 1,4-phenylene, (vi) (nitro)-1,3- or 1,4-phenylene, (vii) (amino)-1,3- or

1,4-phenylene, (viii) (di-lower alkylamino)-1,3- or 1,4-phenylene; and


A_pN-

is monocyclic five to seven ring-membered lower alkyleneamino, piperazino, morpholino, thiamorpholino or N-(lower alkyl, hydroxy, lower alkyl, HPh-lower alkyl or HPh) piperazino;

5 (The most preferred compound of the formula I wherein the RCO is defined in group I_p wherein one of R_{1p} is hydrogen and the R_{2p} is methyl, Ph is -1,4-(3-chlorophenylene) and A_pN- is




(q)



I_q


wherein R_q is hydrogen, halogen, hydroxy, cyano, lower alkyl, hydroxy-lower alkyl, lower alkoxy, acetyl, benzyloxy, lower alkylthio, trifluoromethyl, carboxy, carbo-lower alkoxy, nitro, amino, mono-lower alkylamino, di-lower alkylamino, sulfamoyl, di-lower alkylsulfamoyl or difluoromethylsulfonyl; R_{1q} is halogen, cyano, hydroxy-lower alkyl, lower alkoxy, acetyl, acetamido, benzyloxy, lower alkylthio, trifluoromethyl, hydroxy, carboxy, carbo-lower alkoxy, nitro, amino, mono-lower alkylamino, di-lower alkylamino, sulfamoyl, di-lower alkylsulfamoyl or difluoromethylsulfonyl; or R_q taken together with an adjacent R_{1q} is also lower alkylenedioxy; R_{2q} is



wherein Y_q and X_q , independently, are hydrogen or lower alkyl, and ng is one to seven and R_{3q} is hydrogen, lower alkyl, lower alkoxy-carbonyl-lower alkyl, carboxy-lower alkyl, lower alkanoyl, halo-substituted lower alkanoyl, benzyl, halo-benzyl, benzoyl or halo-benzoyl; and when X_q and Y_q are different, their enantiomers.

40 (The more preferred compounds of the formula I wherein RCO is defined by the group (g) is the compound wherein RCO is the acyl residue of racemic 6-chloro- α -methylcarbazole-2-acetic acid.)

(r)




I_r

55 wherein each of the substituents R^{2r} and R^{3r} is a member of the group consisting of unsubstituted phenyl, naphthyl, thienyl and furyl radicals and phenyl radicals substituted by a substituent selected from the group consisting of halogen, lower alkyl, lower alkoxy, nitro and trifluoromethyl radicals; and wherein R^{1r} is selected from the group consisting of carboxyalkyl- and carboxyalkenyl radicals each containing from two to

five carbons.

(The most preferred compound of the formula I wherein RCO is defined by the group (r) is the compound wherein RCO is the acyl residue of β -(4,5-diphenyloxazol-2-yl)propionic acid.)


5 (s)



15 wherein each of X^{1s} and X^{2s} is a hydrogen atom, a halogen atom, an alkyl group having one to four carbon atoms or an alkoxy group having one to four carbon atoms; each of R^{1s} and R^{2s} is a hydrogen atom or an alkyl group having one to four carbon atoms; A^s is carbonyl, methylene or alkylidene having two to four carbon atoms; Y_s is -O-; and ring P_s represents a pyridine or pyridine N-oxide ring.

20 (The most preferred compound of the formula I wherein RCO is defined by the group (s) is the compound wherein RCO is the acyl residue of 2-(5H-[1]benzopyrano[2,3-b]pyridin-7-yl)propionic acid.)

(t)



wherein:

35 Ar_tCO is an aroyl substituent the Ar_t function of which is a member selected from the group consisting of 2-thienyl, 5-lower alkyl-2-thienyl, 5-halo-2-thienyl, 2-naphthyl and 3-pyridyl, said Ar_tCO being in the meta- or para-position relative to the acetyl function;


40 either of R_t and R_{1t} is hydrogen, the other being a member selected from the group consisting of hydrogen, halo and lower alkyl, provided that, when said R_t is halo or lower alkyl, then said Ar_tCO is in the aforementioned para-position, and when said R_{1t} is halo or lower alkyl, then said Ar_tCO is in the aforementioned meta-position, and further provided that when said R_t or R_{1t} is halo, then said Ar_t is a member selected from the group consisting of 2-thienyl, 5-lower alkyl-2-thienyl and 5-halo-2-thienyl;

either of R_{2t} and R_{3t} is a member selected from the group consisting of hydrogen, allyl and lower alkyl, the other being a member selected from the group consisting of hydrogen and lower alkyl, provided that, when either of said R_{2t} and R_{3t} is allyl, the other is hydrogen, and when either of said R_{2t} and R_{3t} is lower alkyl, the other is a member selected from the group consisting of hydrogen and lower alkyl;

45 R_{2t} and R_{3t} taken together, is an alkylene bridge attached to the α -carbon of the acetic acid function:



55 wherein n_t is an integer from two to five; and wherein lower alkyl as above employed is a radical having from one to five carbon atoms;




10 wherein

Ar_{tii}CO is an aroyl substituent the Ar_{tii} function of which is a member selected from the group consisting of 2-thienyl, 5-methyl-2-thienyl, 5-chloro-2-thienyl, 2-naphthyl and 3-pyridyl, said Ar_{tii}CO being in the meta- or para-position relative to the acetic acid function;


15 either of R_{tii} and R_{1tii} is hydrogen, the other being a member selected from the group consisting of hydrogen, chloro and methyl, provided that, when said R_{tii} is chloro or methyl, then said Ar_{tii}CO is in the aforementioned para-position, and when said R_{1tii} is chloro or methyl, then said Ar_{tii}CO is in the aforementioned meta-position, and further provided that, when said R_{tii} and R_{1tii} is chloro, then said Ar_{tii} is a member selected from the group consisting of 2-thienyl, 5-methyl-2-thienyl and 5-chloro-2-thienyl;

20 either of R_{2tii} and R_{3tii} is a member selected from the group consisting of hydrogen, allyl and lower alkyl, the other being a member selected from the group consisting of hydrogen and lower alkyl, provided that, when either of said R_{2tii} and R_{3tii} is allyl, the other is hydrogen, and when either of said R_{2tii} and R_{3tii} is lower alkyl, the other is a member selected from the group consisting of hydrogen and lower alkyl;

R_{2tii} and R_{3tii}, taken together, is an alkylene bridge attached to the α-carbon of the acetic acid function:



wherein n_{tii} is an integer from two to five; and wherein lower alkyl as above employed is a radical having from one to five carbons;




45 wherein:

Ar_{tii}CO is an aroyl substituent the Ar_{tii} function of which is a member selected from the group consisting of 2-thienyl, 5-lower alkyl-2-thienyl and 5-halo-2-thienyl, said Ar_{tii}CO being in the meta- or para- position relative to the acetic acid function;


50 either of R_{tii} and R_{1tii} is hydrogen, the other being a member selected from the group consisting of hydrogen, halo and lower alkyl, provided that, when said R_{tii} is halo or lower alkyl, then said Ar_{tii}CO is in the aforementioned para-position, and when said R_{1tii} is halo or lower alkyl, then said Ar_{tii}CO is in the aforementioned meta-position;

55 either of R_{2tii} and R_{3tii} is a member selected from the group consisting of hydrogen, allyl and lower alkyl, the other being a member selected from the group consisting of hydrogen and lower alkyl, provided that, when either of said R_{2tii} and R_{3tii} is allyl, the other is hydrogen, and when said R_{2tii} and R_{3tii} is lower alkyl, the other is a member selected from the group consisting of hydrogen and lower alkyl;

R_{2tii} and R_{3tii} together is also an alkylene bridge with the α-carbon of the acetic acid function:




wherein n_{tii} is an integer from two to five; and wherein lower alkyl as above employed is a radical having from one to five carbon atoms; or



wherein:

$\text{Ar}_{\text{tiii}}\text{CO}$ is an aroyl substituent the Ar_{tiii} function of which is a member selected from the group consisting of 2-thienyl, 5-methyl-2-thienyl and 5-chloro-2-thienyl, said $\text{Ar}_{\text{tiii}}\text{CO}$ being in the meta- or para- position relative to the acetic acid function; either of R_{tiii} and $\text{R}_{1\text{tiii}}$ is hydrogen, the other being a member selected from the group consisting of hydrogen, chloro and methyl, provided that, when said R_{tiii} is chloro or methyl, then said $\text{Ar}_{\text{tiii}}\text{CO}$ is in the aforementioned para-position, and when said $\text{R}_{1\text{tiii}}$ is chloro or methyl, then said $\text{Ar}_{\text{tiii}}\text{CO}$ is in the aforementioned meta-position;


either of $\text{R}_{2\text{tiii}}$ and $\text{R}_{3\text{tiii}}$ is a member selected from the group consisting of hydrogen, allyl and lower alkyl, the other being a member selected from the group consisting of hydrogen and lower alkyl, provided that, when either of said $\text{R}_{2\text{tiii}}$ and $\text{R}_{3\text{tiii}}$ is allyl, the other is hydrogen, and when either of said $\text{R}_{2\text{tiii}}$ and $\text{R}_{3\text{tiii}}$ is lower alkyl, the other is a member selected from the group consisting of hydrogen and lower alkyl; $\text{R}_{2\text{tiii}}$ and $\text{R}_{3\text{tiii}}$ together is also an alkylene bridge with the α -carbon of the acetic acid function:



wherein n_{tiii} is an integer from two to five; and wherein lower alkyl as above employed is a radical having from one to five carbon atoms.

(The most preferred compound of the formula I wherein RCO is defined by the group (t) is that of I_{t} wherein Ar_{t} is 2-thienyl, R_{t} and $\text{R}_{1\text{t}}$ are hydrogen, $\text{R}_{2\text{t}}$ is hydrogen and $\text{R}_{3\text{t}}$ is methyl.)

(u)




wherein each of R^{1u} and R^{2u} is a hydrogen atom or an alkyl group having one to four carbon atoms; each of X^{1u} and X^{2u} is a hydrogen atom, a halogen atom, an alkyl group having one to four carbon atoms or an

alkoxy group having one to four carbon atoms; and Y_u is CO.

(The most preferred compound of the formula I wherein RCO is defined by the group (u) is that wherein X^{1u} and X^{2u} is hydrogen, R^{1u} is hydrogen and R^{2u} is methyl.)

(v)



I_v

wherein R_{2v} and R_{3v} each are phenyl or phenyl mono- or di-substituted by at least one member selected from the group consisting of alkyl, alkoxy, alkylmercapto, monoalkylamino, dialkylamino or alkanoylamino wherein the alkyl, alkoxy and alkanoyl each are of up to four carbon atoms, F, Cl, Br, I, CF₃, OH, methylenedioxy, NH₂ and NO₂, A_v is $C_{na}H_{2na}$ wherein na is an integer from one to ten inclusive or




and Z_v is O or S.

(The most preferred compound of the formula I wherein RCO is defined by the group I_v wherein Z_v is oxygen, A_v is



R_{2v} and R_{3v} are each parachlorophenyl.)

(w)



I_w


wherein R_{1w} , R_{2w} , R_{3w} and R_{4w} are independently hydrogen, chlorine, fluorine, bromine, lower alkyl, lower alkoxy and R_{2w} may additionally be trifluoromethyl, and R_{5w} is H or lower alkyl with exclusion of R_{1w} , R_{2w} and R_{3w} as hydrogen simultaneously.

(The most preferred compound of the formula I wherein RCO is defined by the group (w) is that wherein R_{4w} , R_{2w} and R_{5w} are hydrogen and R_{3w} and R_{1w} are each ortho-chloro.)
(x) the acyl residue of (\pm)- α -methyl-3-phenyl-7-benzofuranepropionic acid (I_w).

5

(y)

10



15

where X_Y is



20

$R_{\alpha Y}$ and $R_{\beta Y}$ are hydrogen or lower alkyl and R_Y is halo.

(The most preferred compound of formula I wherein RCO is defined by the group (y) is a compound wherein RCO is the acyl residue of 4-keto-4-(3'-chloro-4'-cyclohexyl)phenylbutyric acid.)

25

Appropriate compounds of formula I are useful in the free base form, in the form of base salts where possible, and in the free acid form, or acid in the form of addition salts where possible. The three forms are within the scope of the invention. In practice, use of the salt form amounts to use of the base form. Pharmaceutically acceptable salts within the scope of the invention may be those derived from mineral acids such as hydrochloric acid and sulfuric acid; and organic acids such as ethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, and the like, giving the hydrochloride, sulfamate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate, and the like, respectively or those derived from bases such as suitable organic and inorganic bases. Examples of pharmaceutically acceptable base addition salts with compounds of the present invention include organic bases which are nontoxic and strong enough to form such salts. These organic bases form a class whose limits are readily understood by those skilled in the art. Merely for purposes of illustration, the class may be said to include mono-, di-, and trialkylamines, such as methylamine, dimethylamine, and triethylamine; mono-, di-, or trihydroxyalkylamines such as mono-, di-, and triethanolamine; amino acids such as arginine, and lysine; guanidine; N-methylglucosamine; N-methylglucamine; L-glutamine; N-methylpiperazine; morpholine; ethylenediamine; N-benzylphenethylamine; tris(hydroxymethyl)aminomethane; and the like. (See for example, "Pharmaceutical Salts," J. Pharm. Sci., 66(1):1-19 (1977).)

30

35

40

The addition salts of said basic or acidic compounds are prepared either by dissolving the free base or acid of compound I in aqueous or aqueous alcohol solution or other suitable solvents containing the appropriate acid or base reagent and isolating the salt by evaporating the solution, or by reacting the free base of compound I with an acid as well as reacting compound I having an acid group thereon with a base such that the reactions are in an organic solvent, in which case the salt separates directly or can be obtained by concentration of the solution.

45

The compounds of the invention may contain an asymmetric carbon atom. Thus, the invention includes the individual stereoisomers, and mixtures thereof. The individual isomers may be prepared or isolated by methods known in the art.

50

By virtue of their lipooxygenase inhibitory properties, said compounds and salts find application in the treatment or prophylaxis of any condition where a lipooxygenase inhibitor is indicated, especially spasmogenic and allergic conditions, psoriasis, and as utility in cytoprotection.

By virtue of their cyclooxygenase inhibitory properties, said compounds and salts find application in the treatment or prophylaxis of any condition where a cyclooxygenase inhibitor is indicated, especially pyrexia, pain, and inflammation.

55

By virtue of both their lipooxygenase and cyclooxygenase inhibitory properties, said compounds and salts find application in the treatment or prophylaxis of any condition where a dual lipooxygenase/cyclooxygenase inhibitor is indicated, especially any condition involving blood platelet aggregation or inflammation. In the case of inflammation, the compounds and salts are particularly suited to the treatment or prophylaxis of conditions associated with infiltration of leukocytes into inflamed tissue.

In determining when a lipoxygenase, cyclooxygenase, or dual lipoxygenase/cyclooxygenase inhibitor is indicated, of course *inter alia*, the particular condition in question and its severity, as well as, the age, sex, weight, and the like of the subject to be treated, must be taken into consideration and this determination is ultimately at the discretion of the attendant physician.

6 Examples of the aforesaid spasmogenic conditions are those involving smooth muscle tissue, especially airway smooth muscle constriction such as intrinsic asthma--(including intrinsic or idiopathic bronchial asthma and cardiac asthma), bronchitis and arterial smooth muscle constriction such as coronary spasm (including that associated with myocardial infarction, which may or may not lead to left ventricular failure resulting in cardiac asthma) and cerebral spasm or 'stroke'. Other examples include bowel disease caused
10 by abnormal colonic muscular contraction such as may be termed 'irritable bowel syndrome', 'spastic colon' or 'mucous colitis'.

Examples of the aforesaid allergic conditions are extrinsic asthma (from which it will be appreciated that said compounds and salts are particularly favorable as antiasthmatic agents), allergic skin diseases such as eczema having a total or partial allergic origin, allergic bowel disease (including coeliac disease) and
15 allergic eye conditions, hay fever, and allergic conjunctivitis.

Examples of the aforesaid pyretic and painful conditions include fever associated with infections, trauma and injury, malignant disease, and diseases affecting the immune system (including autoimmune diseases).

Examples of the aforesaid conditions involving blood platelet aggregation are those resulting from thrombosis, including 'stroke' having a total or partial thrombotic origin, coronary thrombosis, phlebitis, and
20 phlebothrombosis (the latter two conditions also possibly being associated with inflammation).

Examples of the aforesaid conditions involving inflammation are inflammatory conditions of the lung, joints, eye, bowel, skin, and heart.

Inflammatory lung conditions which may be so treated and/or prevented include asthma and bronchitis (*vide supra*) and cystic fibrosis (which may also or alternatively involve the bowel or other tissue).

25 Inflammatory joint conditions which may be so treated and/or prevented include rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, gouty arthritis, and other arthritic conditions.

Inflammatory eye conditions which may be so treated and/or prevented include uveitis (including intis) and conjunctivitis (*vide supra*).

Inflammatory bowel conditions which may be so treated and/or prevented include Crohn's disease,
30 ulcerative colitis, and ischemic bowel disease.

Inflammatory skin diseases which may be so treated and/or prevented include those associated with cell proliferation, such as psoriasis and eczema (*vide supra*) and dermatitis (whether or not of allergic origin).

Inflammatory conditions of the heart which may be so treated and/or prevented include coronary infarct
35 damage.

Other inflammatory conditions which may be so treated and/or prevented include tissue necrosis of chronic inflammation and tissue rejection following transplant surgery.

For medical use, the amount required of a compound of formula (I) or physiologically acceptable salt thereof--(hereinafter referred to as the active ingredient) to achieve a therapeutic effect will, of course, vary
40 both with the particular compound, the route of administration and the mammal under treatment and the particular disorder or disease concerned. A suitable dose of a compound of formula (I) or physiologically acceptable salt thereof for a mammal suffering from, or likely to suffer from any condition as described hereinbefore is 0.1 μ g-500 mg of base per kilogram body weight. In the case of systemic administration, the dose may be in the range 0.5 to 500 mg of base per kilogram body weight, the most preferred dosage
45 being 0.5 to 50 mg/kg of mammal body weight for Example 5 to 25 mg/kg; administered two or three times daily. In the case of topical administration, e.g. to the skin or eye, a suitable dose may be in the range 0.1 ng-100 μ g of base per kilogram, typically about 0.1 μ g/kg.

In the case of oral dosing for the treatment or prophylaxis of airway smooth muscle constriction, or asthma, or bronchitis in general, due to any cause, a suitable dose of a compound of formula (I) or
50 physiologically acceptable salt thereof, may be as specified in the preceding paragraph, but most preferably is from 1 mg to 10 mg of base per kilogram, the most preferred dosage being from 1 mg to 5 mg/kg of mammal body weight, for example from 1 to 2 mg/kg. In the case of pulmonary administration for the latter indications, the dose may be in the range of from 2 μ g to 100 mg, for example from 20 μ g to 0.5 mg, especially 0.1 to 0.7 mg/kg.

55 It is understood that the ordinarily skilled physician or veterinarian will readily determine and prescribe the effective amount of the compound to prevent or arrest the progress of the condition for which treatment is administered. In so proceeding, the physician or veterinarian could employ relatively low doses at first, subsequently increasing the dose until a maximum response is obtained.

While it is possible for an active ingredient to be administered alone, it is preferable to present it as a pharmaceutical formulation comprising a compound of formula (I) or a pharmacologically acceptable acid addition salt thereof and a physiologically acceptable acid addition salt thereof and a physiologically acceptable carrier therefor. Such formulations constitute a further feature of the present invention. Conveniently, the active ingredient comprises from 0.1% to 99.9% by weight of the formulation. Conveniently, unit doses of a formulation contain between 0.1 mg and 1 g of the active ingredient. For topical administration, the active ingredient preferably comprises from 1% to 2% by weight of the formulation but the active ingredient may comprise as much as 10% w/w. Formulations suitable for nasal or buccal administration, (such as self-propelling powder dispensing formulations described hereinafter), may comprise 0.1 to 20% w/w, for example 2% w/w of active ingredient.

The formulations, both for veterinary and for human medical use, of the present invention comprise an active ingredient in association with a pharmaceutically acceptable carrier therefor and optionally other therapeutic ingredient(s). The carrier(s) must be 'acceptable' in the sense of being compatible with the other ingredients of the formulations and not deleterious to the recipient thereof.

The formulations include those in a form suitable for oral, pulmonary, ophthalmic, rectal, parenteral (including subcutaneous, intramuscular, and intravenous), intraarticular, topical, nasal, or buccal administration.

The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well-known in the art of pharmacy. All methods include the step of bringing the active ingredient into association with the carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing the active ingredient into association with a liquid carrier or a finely divided solid carrier or both, and then, if necessary, shaping the product into the desired formulation.

Formulations of the present invention suitable for oral administration may be in the form of discrete units such as capsules, cachets, tablets, or lozenges, each containing a predetermined amount of the active ingredient; in the form of a powder or granules; in the form of a solution or a suspension in an aqueous liquid or nonaqueous liquid; or in the form of an oil-in-water emulsion or a water-in-oil emulsion. The active ingredient may also be in the form of a bolus, electuary, or paste.

A tablet may be made by compressing or molding the active ingredient optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing, in a suitable machine, the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, surface active, or dispersing agent. Molded tablets may be made by molding, in a suitable machine, a mixture of the powdered active ingredient and a suitable carrier moistened with an inert liquid diluent.

Formulations for rectal administration may be in the form of a suppository incorporating the active ingredient and a carrier such as cocoa butter, or in the form of an enema.

Formulations suitable for parenteral administration conveniently comprise a sterile aqueous preparation of the active ingredient which is preferably isotonic with the blood of the recipient.

Formulations suitable for intraarticular administration may be in the form of a sterile aqueous preparation of the active ingredient which may be in microcrystalline form, for example, in the form of an aqueous microcrystalline suspension. Liposomal formulations or biodegradable polymer systems may also be used to present the active ingredient for both intraarticular and ophthalmic administration.

Formulations suitable for topical administration include liquid or semi-liquid preparations such as liniments, lotions, applications, oil-in-water or water-in-oil emulsions such as creams, ointments or pastes; or solutions or suspensions such as drops. For example, for ophthalmic administration, the active ingredient may be presented in the form of aqueous eye drops as, for example, a 0.1-1.0% solution. Formulations suitable for administration to the nose or buccal cavity include powder, self-propelling and spray formulations such as aerosols and atomizers. The formulations, when dispersed, preferably have a particle size in the range of 0.1 to 200 μ .

A particularly valuable form of a pharmaceutical composition of the present invention, for use in the prophylaxis or treatment of airway smooth muscle constriction, or asthma or bronchitis in general, due to any cause, is one suitable for pulmonary administration via the buccal cavity. Preferably the composition is such that particles having a diameter of 0.5 to 7 μ , most preferably 1 to 6 μ , containing active ingredient, are delivered into the lungs of a patient. Such compositions are conveniently in the form of dry powders for administration from a powder inhalation device or self-propelling powder-dispensing containers, for example as a self-propelling aerosol composition in a sealed container; preferably the powders comprise particles containing active ingredient of which particles at least 98% by weight have a diameter greater than 0.5 μ and at least 95% by number have a diameter less than 7 μ . Most desirably at least 95% by weight of the

particles have a diameter greater than 1 μ at least 90% by number of the particles have a diameter less than 6 μ .

The compositions in the form of dry powders preferably include a solid fine powder diluent such as sugar and are conveniently presented in a permeable capsule, for example of gelatin.

5 Self-propelling compositions of the invention may be either powder-dispensing compositions or compositions dispensing the active ingredient in the form of droplets of a solution or suspension. Self-propelling powder-dispensing compositions include a liquid propellant having a boiling point of below 65° F at atmospheric pressure. Generally the propellant may constitute 50 to 99.9% w/w of the composition whilst the active ingredient may constitute 0.1 to 20% w/w, for example about 2% w/w of the composition. The
10 carrier in such compositions may include other constituents, in particular a liquid nonionic or solid anionic surfactant, or a solid diluent (preferably having a particle size of the same order as of the particles of active ingredient) or both. The surfactant may constitute from 0.01 up to 20% w/w, though preferably it constitutes below 1% w/w of the composition.

Self-propelling compositions wherein the active ingredient is present in solution comprise an active
15 ingredient, propellant, and co-solvent, and advantageously an antioxidant stabilizer. The co-solvents may constitute 5 to 40% w/w of the composition, though preferably less than 20% w/w of the composition.

Compositions of the present invention may also be in the form of aqueous or dilute alcoholic solution, optionally a sterile solution, of the active ingredient for use in a nebulizer or atomizer.

Formulations of the present invention may also be in the form of an aqueous or dilute alcoholic solution,
20 optionally a sterile solution, of the active ingredient for use in a nebulizer or atomizer, wherein an accelerated air stream is used to produce a fine mist consisting of small droplets of the solution. Such formulations usually contain a flavoring agent such as saccharin sodium and a volatile oil. A buffering agent and a surface active agent may also be included in such a formulation which should also contain a preservative such as methylhydroxybenzoate.

25 Other formulations suitable for nasal administration include a coarse powder having a particle size of 20 to 500 microns which is administered in the manner in which snuff is taken i.e. by rapid inhalation through the nasal passage from a container of the powder held close up to the nose.

In addition to the aforementioned ingredients, the formulations of this invention may include one or
more additional ingredients such as diluents, buffers, flavoring agents, binders, surface active agents,
30 thickeners, lubricants, preservatives e.g. methylhydroxybenzoate (including antioxidants), emulsifying agents, and the like. Any other therapeutic ingredient may comprise one or more of the following: antibiotic (e.g. antibacterial), antifungal and antiviral agents, and antihistamines (particularly peripherally acting antihistamines). However, when such other agent(s) are also present, according to another aspect of the invention, the compound of formula (I) or physiologically acceptable salt thereof and the other agent(s),
35 need not necessarily be present as a pharmaceutical formulation as hereinbefore defined, but merely in combination or intimate admixture, i.e. optionally, a pharmaceutically acceptable carrier need not be present.

The combination with antihistamines is particularly favored for antiasthmatic use. Such an antihistamine may be selected from any compound described in European Patent Applications EP O 859 949 A and EP
40 O 117 302 A. The amount and dosage regime for such an antihistamine may be chosen from any of those recited in the latter two European Specifications. Especially preferred are the antihistamines (E)-3-(6-(3-pyrrolidino)-1-(4-tolyl)prop-1E-enyl(-2-pyridyl)) acrylic acid and (E)-3-(6-(3-pyrrolidino)-1-(4-tolyl)prop-1E-enyl(-2-pyridyl))propionic acid. Another preferred antihistamine is (E)-1-(4-methylphenyl)-1-(2-pyridyl)-3-pyrrolidinoprop-1-ene, otherwise known as typrolidine.

45 Also preferred is the antihistamine known as Seldane.

The compound of formula I and their salts may be prepared generally by one of two of the following processes and constitute a further aspect of the present invention.

For one of the preparations of compounds of the formula I as defined above, a compound of the formula

50 RCOOH

wherein R is as defined above;


is treated with a chloro- or bromo-generating compound, such as SOCl_2 , $(\text{COCl})_2$, PBr_3 , preferably SOCl_2 or $(\text{COCl})_2$ under conditions analogous to those known in the art to obtain a compound of the formula

RCOQ

55 wherein Q is chloro or bromo and R is as defined above, and then

RCOQ

is treated with a compound of the formula



5

or its acid addition salt in the presence of a base to give the compound of the formula I. (See Scheme I)

Where RCO is an acyl residue having a sulfinyl substituent, such as in I_b, or an acyl residue having the definition of the formula of I_w then the preparation with treatment by a chloro- or bromo- generating compound, such as SOCl₂, (COCl)₂, or PBr₃ described above does not provide a desired product. For this definition of RCO, an alternate preparation is thus required.

That is, surprisingly, for compounds having RCO defined by I_b having a sulfoxide group among the definitions of its substituents, the sulfoxide reacts with the chloro- or bromo- of the generating compound so a product of the formula I having the sulfoxide as a substituent cannot be obtained. In other words, the acyl residue of sulindac does not provide the RCON(OR₁)R₂ desired by use of the above desired procedure.

Further, in like manner, the compounds having the formula RCO of the structure I_w cannot be treated with a chloro- or bromo- generating compound, such as SOCl₂, (COCl)₂, or PBr₃ to obtain the desired product of the above preparation because an intramolecular cyclization to oxindole occurs. For example, the acyl residue of methyl diclofenate as shown in U.S. Patent No. 4,092,430 cannot be treated as described in the above preparation and reacted with the substituted hydroxyamine hydrogen chloride of the present invention contrary to the teaching of U.S. Patent No. 4,092,430.

Therefore, in these two instances use of the alternate preparation described hereinafter is critical.

In an alternate preparation of the compound of the formula I as defined above a compound of the formula RCOOH

is treated with a coupling agent, such as 1,1'-carbonyldiimidazole, N-ethoxycarbonyl-2-ethoxy-1,3-dihydroquinoline, dicyclohexylcarbodiimide, dicyclohexylcarbodiimide/hydroxybenzotriazole or the like, preferably 1,1'-carbonyldiimidazole; and then further treated with a compound of the formula

30



35

or the acid addition salt thereof; wherein R₁ and R₂ are as defined above; to obtain the compound of formula I. (See Scheme II)


40

45


50

55

Scheme I



Scheme II



The reactions of Scheme II are carried out in organic solvents such as methylene chloride, DMF, THF and the like, preferably in methylene chloride.

The hydroxamic acid formation is carried out at a temperature ranging from about 0° C to about 50° C, preferably at about room temperature.

One of skill in the art would recognize variations in the sequence and would recognize appropriate reaction conditions from analogous reactions which may be appropriately used in the processes to make the compounds of formula (I) herein. Further, the starting materials are known or can be prepared by known methods.

Under certain circumstances it is necessary to protect either the N or O of intermediates in the above noted process with suitable protecting groups which are known. Introduction and removal of such suitable oxygen and nitrogen protecting groups are well-known in the art of organic chemistry; see for example "Protective Groups in Organic Synthesis," T. W. Greene, (Wiley-Interscience), New York, 1981.

Examples of suitable oxygen protecting groups are benzyl, t-butyldimethylsilyl, ethoxyethyl, and the like. Protection of an N-H containing moiety is necessary for some of the processes described herein for the preparation of compounds of this invention. Suitable nitrogen protecting groups are benzyl, triphenylmethyl, trialkylsilyl, trichloroethylcarbamate, trichloroethoxycarbonyl, vinyloxycarbamate, and the like.

Under certain circumstances it is necessary to protect two different oxygens with dissimilar protecting groups such that one can be selectively removed while leaving the other in place. The benzyl and t-butyldimethylsilyl groups are used in this way; either is removable in the presence of the other, benzyl being removed by catalytic hydrogenolysis, and t-butyldimethylsilyl being removed by reaction with, for example, tetra-n-butylammonium fluoride.

In the process described herein for the preparation of compounds of this invention the requirements for protective groups are generally well recognized by one skilled in the art of organic chemistry, and accordingly the use of appropriate protecting groups is necessarily implied by the processes of the charts herein, although not expressly illustrated.

The products of the reactions described herein are isolated by conventional means such as extraction, column chromatography, flash chromatography, and the like.

The salts of the compounds of formula (I) described above are prepared by reacting the appropriate base or acid with a stoichiometric equivalent of the compounds of formula (I), respectively, to obtain

pharmaceutically acceptable salts thereof.

The invention is further elaborated by the representative examples as follows. Such examples are not meant to be limiting.

EXAMPLES

EXAMPLE 1

(Z)-6-fluoro-N-hydroxy-N,2-dimethyl-3-[[4-(methylsulfinyl)phenyl]methylene]-3H-indene-1-acetamide.

A mixture of (Z)-5-fluoro-2-methyl-1-[[4-(methylsulfinyl)phenyl]methylene]-1H-indene-3-acetic acid (sulindac, Sigma) (7.13 g, 0.02 mole) and 1,1'-carbonyldiimidazole (4.9 g, 0.03 mole) in methylene chloride (200 ml) is stirred at room temperature for 30 minutes, then N-methyl-hydroxylamine hydrochloride (3.34 g, 0.04 mole) is added. After being stirred an additional 41 hours, the mixture is poured into cold 2 normal hydrochloric acid (1 l) and extracted with methylene chloride (400 ml). The organic solution is washed successively with 2 normal hydrochloric acid (1 l), water (3 x 1 l) and is dried over sodium sulfate. The solvent is removed under reduced pressure and the residue (8.2 g) is chromatographed on silica gel (225 g). Elution with 95:5 mixture of chloroform and methanol, gives 7.35 g of a solid. Recrystallization from ethyl acetate-hexane, then from methylene chloride-hexane, gives 5.5 g (75.3%) of analytically pure (Z)-6-fluoro-N-hydroxy-N,2-dimethyl-3-[[4-(methylsulfinyl)phenyl]methylene]-3H-indene-1-acetamide, m.p. 156-160 °C.

EXAMPLE 2

(Z)-6-fluoro-N-methoxy-2-methyl-3-[[4-(methylsulfinyl)phenyl]methylene]-3H-indene-1-acetamide.

Prepared by the method described in Example 1, using appropriate starting materials. Recrystallization from methanol-ethyl acetate-hexane, gives analytically pure (Z)-6-fluoro-N-methoxy-2-methyl-3-[[4-(methylsulfinyl)phenyl]methylene]-3H-indene-1-acetamide in 50.8% yield, m.p. 177-179 °C.

EXAMPLE 2A

(Z)-6-fluoro-N-hydroxy-2-methyl-3-[[4-(methylsulfinyl)phenyl]methylene]-3H-indene-1-acetamide.

Prepared by the method described in Example 1, using appropriate starting materials. Recrystallization from methanol/ethyl acetate/hexane gives analytically pure (Z)-6-fluoro-N-hydroxy-2-methyl-3-[[4-(methylsulfinyl)phenyl]methylene]-3H-indene-1-acetamide 28.9% yield, m.p. 208-209 °C dec.

EXAMPLE 2B

(Z)-6-Fluoro-N-hydroxy-N-(1-methylethyl)-2-methyl-3-[[4-(methylsulfinyl)phenyl]methylene]-3H-indene-1-acetamide.

Prepared by the method described in Example 2, using appropriate starting materials. The crude product is flash chromatographed over silica gel using 5% CH₃OH-CHCl₃ as eluent and obtained the pure

analytical product as viscous oil in 94.8% yield.

EXAMPLE 3

5

1-(4-Chlorobenzoyl)-N-hydroxy-5-methoxy-N,2-dimethyl-1H-indole-3-acetamide.

10 Oxalyl chloride (56.92 g, 0.448 mole) is added dropwise over 20 minutes with stirring to a solution of 1-(p-chlorobenzoyl)-5-methoxy-2-methylindole-3-acetic acid (indomethacin, sigma) (64.2 g, 0.179 mole) in methylene chloride (500 ml) and dimethylformamide (13.12 g, 0.179 mole) at $< -2^{\circ}\text{C}$. After being stirred for 80 minutes, this solution is added to a solution of N-methylhydroxylamine hydrochloride (59.9 g, 0.717 mole) and triethylamine (163.3 g, 1.61 mole) in tetrahydrofuran (200 ml) and water (100 ml). After being
15 stirred for an additional 75 minutes, the mixture is poured into cold 2 normal hydrochloric acid (3 l) and extracted with methylene chloride (1.5 l). The organic layer is washed with water and dried over sodium sulfate. The solvent is removed under reduced pressure and the residue (68.7 g) is chromatographed on 663 g of silica gel. Elution with chloroform, then with 95:5 mixture of chloroform-methanol gives 63.1 g of a solid. Recrystallization from methanol, gives 55.49 g (80%) of analytically pure 1-(4-chlorobenzoyl)-N-
20 hydroxy-5-methoxy-N,2-dimethyl-1H-indole-3-acetamide, m.p. $170-171^{\circ}\text{C}$.

EXAMPLE 4

25

1-(4-Chlorobenzoyl)-N,5-dimethoxy-2-methyl-1H-indole-3-acetamide.

Prepared by the method described in Example 3 using appropriate starting materials. Recrystallization
30 from methanol, gives analytically pure 1-(4-chlorobenzoyl)-N,5-dimethoxy-2-methyl-1H-indole-3-acetamide in 73.7% yield, m.p. $184-185^{\circ}\text{C}$.

EXAMPLE 5

35

1-(4-Chlorobenzoyl)-N-hydroxy-5-methoxy-2-methyl-N-(1-methylethyl)-1H-indole-3-acetamide.

40 Prepared by the method described in Example 3 using appropriate starting materials. The crude product is chromatographed over silica gel using chloroform as eluant. Recrystallization from methanol gives analytically pure 1-(4-chlorobenzoyl)-N-hydroxy-5-methoxy-2-methyl-N-(1-methylethyl)-1H-indole-3-acetamide in 41.6% yield, m.p. $198-199^{\circ}\text{C}$.

45

EXAMPLE 6

1-(4-Chlorobenzoyl)-N-cyclohexyl-N-hydroxy-5-methoxy-2-methyl-1H-indole-3-acetamide.

Prepared by the method described in Example 3 using appropriate starting materials. The crude product is chromatographed on silica gel using chloroform as eluant. Recrystallization from methanol gives
55 analytically pure 1-(4-chlorobenzoyl)-N-cyclohexyl-N-hydroxy-5-methoxy-2-methyl-1H-indole-3-acetamide in 49.2% yield, m.p. $208-210^{\circ}\text{C}$.

EXAMPLE 6A

1-(4-Chlorobenzoyl)-N,5-dimethoxy-N,2-dimethyl-1H-indole-3-acetamide.

Prepared by the method described in Example 3 using appropriate starting materials. The crude product is flash chromatographed over silica gel using CHCl_3 as eluent. Recrystallization from CH_2Cl_2 -isoPr₂O gives analytically pure product in 86.5% yield, m.p. 155-157 °C.

EXAMPLE 6B

1-(4-Chlorobenzoyl)-N-hydroxy-5-methoxy-2-methyl-N-phenyl-1H-indole-3-acetamide.

Prepared by the method described in Example 3 using appropriate starting materials. The crude product is flash chromatographed over silica gel using CHCl_3 as eluent. Recrystallization of the purified product from methanol gives analytically pure product in 11.4% yield, m.p. 173-174 °C.

EXAMPLE 6C

1-(4-Chlorobenzoyl)-N-benzyl-N-hydroxy-5-methoxy-2-methyl-1H-indole-3-acetamide.

Prepared by the method described in Example 3 using appropriate starting materials. The crude product is flash chromatographed over silica gel using 5% $\text{CH}_3\text{OH}-\text{CH}_2\text{Cl}_2$ as eluent. Recrystallization of the purified product from ethylacetate-ether gives analytically pure product in 27.6% yield, m.p. 188-189 °C.

EXAMPLE 7

N-Methoxy-2-[(2,6-dichlorophenyl)amino]-benzeneacetamide.

A warm solution of 2-[(2,6-dichlorophenyl)amino]benzeneacetic acid (2.03 g; 7 mmol) in CH_2Cl_2 (100 mL) is treated with 1,1'-carbonyl-diimidazole (1.75 g; 11 mmol) and the mixture is stirred at room temperature for 5.0 minutes when a clear solution is obtained. The solution is then treated with methoxyamine hydrochloride (1.50 g; 18 mmol) and the reaction mixture stirred at room temperature for 108 hours. It is then decomposed with water, extracted with CH_2Cl_2 , dried over Na and then evaporated to dryness to give a solid. Fractional recrystallization of the crude product from a mixture of toluene- CH_2Cl_2 gives off-white crystalline solid (1.27 g; 55.8%); mp 177-178 °C.

EXAMPLE 8

N-Hydroxy-N-methyl-2-[(2,6-dichlorophenyl)amino]benzeneacetamide.

1,1'-Carbonyl-diimidazole (5.5 g; 34 mmol) is slowly added to a warm solution of 2-[(2,6-dichlorophenyl)amino]benzeneacetic acid (6.5 g; 22 mmol) in CH_2Cl_2 (600 mL) and stirred at room temperature for 2.0 hours when a clear solution is formed. N-Methylhydroxylamine hydrochloride (5.5 g; 66 mmol) is slowly added to the solution and the mixture stirred at room temperature for 48 hours. It is decomposed with water, extracted with CH_2Cl_2 and dried over Na_2SO_4 . The solvent is evaporated off and the residue

flash chromatographed over silica gel (250 g) using 10% EtOH-CH₂Cl₂ as eluant to give the pure product as white solid which is recrystallized from toluene-CH₂Cl₂. Yield 2.95 g (41.2%); mp 135-138 °C.

The usefulness of the compounds of the present invention as inhibitors of the 5-lipoxygenase enzyme, cyclooxygenase, or other related biochemical actions may be demonstrated by their effectiveness in various standard test procedures. A description of each procedure follows.

ARBL/ARBC Whole Cell 5-Lipoxygenase and Cyclooxygenase Assays

10

Materials

The rat basophilic leukemia cell line (RBL-1) was obtained from the American Type Culture Collection (Rockville, MD).

Radioimmuno assay (RIA) kits of LTB₄ and PGF_{2α} were obtained from Amersham (Arlington Heights, IL) and Seragen (Boston, MA) respectively.

All tissue culture media were obtained from GIBCO (Grand Island, NY).

20

Method

RBL-1 cells are grown in suspension culture in Eagle's minimum essential medium supplemented with 12% fetal bovine serum at 37 °C in an incubator supplied with air-5% carbon dioxide. Cells are harvested by centrifugation. They are washed with cold phosphate buffered saline pH 7.4 (PBS; NaCl, 7.1 g; Na₂HPO₄ 1.15 g; KH₂PO₄, 0.2 g; and KCl, 0.2 g/l). Cells are finally suspended in PBS containing 1.0 mM calcium at a density of 2x10⁶ cells/ml. Cells are incubated with and without test agent (in DMSO) (1% DMSO is without effect on arachidonic acid metabolism) for ten minutes at room temperature. Calcium ionophore A23187 (5 μM) is added and cells are incubated for seven minutes at 37 °C. The reaction is stopped by chilling the tubes on ice for ten minutes. Cells are separated by centrifugation and the supernatant is stored at -20 °C. Aliquots (100 μl) are analyzed for LTB₄ and PGF_{2α} using radioimmuno assay kits as provided by the supplier.

Table 1 contains biochemical data obtained from this whole cell assay as IC₅₀s which are calculated as the amount of test compound causing 50% inhibition of LTB₄ or PGF_{2α} formation.

35

Carrageenan-Induced Rat Foot Paw Edema-2 (CFE-2) Assay:

40

Protocol

Carrageenan solution (1% w/v) is prepared by dissolving 100 mg carrageenan (Marine Colloidal DIV., Springfield, NJ) in 10 ml of sterile saline (0.9%) solution (Travenol). The solution is vortexed for thirty to forty-five minutes. Animals are dosed with compound one hour before carrageenan challenge. Foot paw edema is induced by injecting 0.05 ml of the 1% carrageenan subcutaneously into the plantar portion of the right hind paw of each rat under light anesthesia. Initial foot paw volume is measured immediately following carrageenan challenge using mercury plethysmography (Buxco Electronics). Edema is measured five hours after carrageenan. The difference between the five-hour and the initial paw volume is expressed as delta edema. The delta edema for each test group of animals is used to calculate the percent inhibition of edema achieved by the compound at the test dose compared with the vehicle control group. The ID₂₅ (the dose at which swelling is inhibited by 25%) was calculated by probit analysis for the dose at which a result of 25 percent inhibition occurs.

55

Mycobacterium - Induced Rat Footpad Edema Assay (MFE):

Protocol

5 Myobacterium butyricum (5 mg/ml) is suspended in paraffin oil by sonication for ten minutes in an ice bath. Footpad edema is induced on Day 0 by injecting 0.1 ml of the Mycobacterium mixture into the left hindpaw of lightly anesthetized rats. Swelling in the injected hindpaw is determined by mercury plethysmography seventy-two hours after injection. Groups of rats are treated with test compounds (suspended in 0.5% hydroxypropyl methylcellulose with 0.2% Tween-80) or vehicle one hour before Myobacterium injection and on Days 1 and 2. Inhibition of swelling is determined by comparing the change in hindpaw
10 volume in compound- and vehicle-treated rats. An ID_{40} (the dose at which swelling is inhibited by 40%) was calculated by probit analysis.

Gastric Ulcerogenicity (UD)

15

Protocol

20

Male outbred Wistar rats (100-250 gms) were fasted for twenty-four hours. After fasting, test compounds were administered orally (in 2 ml/Kg of 0.5% hydroxypropyl methylcellulose) and the rats were denied access to food and water for six more hours. The rats were then sacrificed with CO_2 so that the stomachs could be removed, opened along the greater curvature, and evaluated for the presence of gastric
25 ulcers. Results are expressed as the percent of rats with gastric ulcers at a given dose.

The results of the CFE-2, MFE, and UD assays for each of the noted compounds is shown in the following Table 1 and Table 2.

30

35

40

45

50

55

TABLE I
SULINDAC SERIES

PD #	X	ARBC ^a /ARBL ^b	RATIO ^c	CFE ^d	MFE ^e
SULINDAC					
5 Example 1	CON(ME)OH	0% @ 32*/1.0	---	15% @ 30	30% @ 10
Example 2	CONHOME	0% @ 32*/26.4	---	---	---
Example 2A	CONHOH	0% @ 32*/12.9	---	31% @ 30	53% @ 50

*Parent sulindac sulfoxides are not CO inhibitors, but converted to CO Inhibitory sulfides in vivo.

^a ARBC = RBL Whole Cell Cyclooxygenase Assay. IC50 (μM) for PGF_{2α} inhibition.

^b ARBL = RBL Whole Cell 5-Lipoxygenase Assay. IC50 (μM) for LTB₄ inhibition.

^c RATIO = Selectivity of Inhibition. If > 1.0 compound is cyclooxygenase selective.
If < 1.0 compound is 5-lipoxygenase selective.

^d CFE = Carrageenan Footpad Edema Model. ID25 or % Inhibition @ Dose (mg/kg PO).

^e MFE = Mycobacterium Footpad Edema Model. ID40 or % Inhibition @ Dose (mg/kg PO).

TABLE 2
INDOMETHACIN SERIES

Example #	X	ARBC ^a /ARBL ^b	RATIO ^c	CFE ^d	MFE ^e	UD50 ^f
Indomethacin	OH	0.5/ > 100	> 200	36% @ 5	0.21	5.4
Oxametacin	NHOH	1.1/7.5	6.8	46% @ 30		103
Example 3	NMEOH	5.2/1.4	0.27	29% @ 10	< 2	10% @ 100
Example 4	NHOME	0.2/24	120	40% @ 10		42
Example 5	N(iPR)OH	2.7/0.9	0.33	27% @ 10	0.82	N @ 100
Example 6	N(C-C ₆ H ₁₁)OH	0.1/1.6	16.0	22% @ 10	4.7	N @ 200

^a ARBC = RBL Whole Cell Cyclooxygenase Assay. IC50 (μM).

^b ARBL = RBL Whole Cell 5-Lipoxygenase Assay. IC50 (μM).

^c RATIO = Selectivity of Inhibition. If > 1:0 compound is cyclooxygenase selective.
If < 1:0 compound is 5-lipoxygenase selective.

^d CFE = Carrageenan Footpad Edema Model. ID25 or % Inhibition @ Dose (mg/kg PO).

^e MFE = Mycobacterium Footpad Edema Model. ID40 or % Inhibition @ Dose (mg/kg PO).

^f UD50 = Dose (mg/kg PO) causing ulcers in 50% of animals.

Accordingly, the present invention is

- (a) a compound of formula I or an acid addition salt thereof;
- (b) a method for preparing a compound of formula I or a pharmacologically acceptable acid addition salt thereof;
- (c) a pharmaceutical formulation comprising a compound of formula (I) or a physiologically acceptable salt thereof and a pharmaceutically acceptable carrier therefor;
- (d) a method for preparing such formulations for the inhibition of the lipoxigenase and/or cyclooxygenase pathways of the arachidonic acid metabolism by use of a nontoxic, effective, inhibitory amount of a compound of formula I or a physiologically acceptable salt thereof;
- (e) a method for preparing such formulations for the prophylaxis or treatment of disease or condition in a mammal, including man, comprising the administration to said mammal of a nontoxic, therapeutically or prophylactically effective amount of a compound of formula I or a physiologically acceptable salt thereof;
- (f) a method for preparing such formulations for the prophylaxis or treatment of any individual condition described herein, in a mammal, including man, comprising the administration to said mammal of a nontoxic therapeutically or prophylactically effective amount of a compound of formula I or a physiologically acceptable salt thereof;
- (g) a method for preparing such formulations for the prophylaxis or treatment of inflammation, arthritis, pain or fever in a mammal, including man, comprising the administration to said mammal of a nontoxic, effective, antiinflammation, antiarthritic, analgesic, antipyretic amount of a compound of formula I or a physiologically acceptable salt thereof;
- (h) a method for preparing such formulations for the prophylaxis or treatment of asthma in a mammal, including man, comprising administration to said mammal of a nontoxic, effective, antiasthmatic amount of a compound of formula I or a physiologically acceptable salt thereof;
- (i) a compound of formula I or a physiologically acceptable salt thereof for use in medicine, especially as defined in (d)-(h) above;
- (j) use of a compound of formula I or a physiologically acceptable salt thereof in the manufacture of medical therapeutic agents, particularly those for use as defined in (d)-(i) above; and
- (k) any novel feature described herein.


Claims

1. A compound of the formula




and pharmaceutically acceptable acid addition or base salts thereof wherein

- i) RCO is an acyl residue of selected NSAIDS;
- ii) R₁ is hydrogen, lower alkyl or acyl;
- iii) R₂ is H, lower alkyl, cycloalkyl of from three to twelve carbon atoms of which from three to seven carbons are ring carbons, aryl, arylalkyl or heteroaryl with the overall proviso that when the acyl residue is of the formula




wherein R_4 is hydrogen or lower alkyl, then R_3 is hydrogen, phenylmethyl, allyl, vinyl, isopropenyl, benzoyl, chlorobenzoyl, methoxybenzoyl, and thiomethylbenzoyl; and when the acyl residue is of the formula



wherein R_5 , R_6 , R_7 and R_8 are independently H, chlorine, fluorine, or bromine, lower alkyl, lower alkoxy, or trifluoromethyl but when R_4 is hydrogen then are not all hydrogen; and R_4 is otherwise independently as defined above; then R_1 and R_2 cannot both be hydrogen.

2. A compound of Claim 1 wherein RCO is selected from the groups consisting of

(a)



in which

R_{1a} is selected from the group consisting of benzene, naphthalene, biphenyl and substituted benzene, naphthalene and biphenyl radicals in which said substituent is selected from the group consisting of halogen, lower alkyl, lower alkylthio, lower alkoxy, trifluoromethyl, phenoxy, lower alkyl phenoxy, lower alkoxy phenoxy, halogenophenoxy, trifluoroacetyl, difluoroacetyl, monofluoroacetyl, di-lower alkyl sulfamyl, lower alkanoyl, di-lower alkyl carboxamido, cyano, carb-lower alkoxy, trifluoromethylthio, lower alkyl sulfinyl, lower alkylsulfonyl, benzylthio, lower alkylbenzylthio, lower alkoxybenzylthio, halogenobenzylthio, mercapto, nitro, amino, di-(lower alkyl)amino, lower alkylamino, lower alkanoylamino, hydroxy, lower alkanoyloxy, trifluoroacetoxy, difluoroacetoxy, monofluoroacetoxy, benzyloxy, lower alkylbenzyloxy, lower alkoxybenzyloxy, and halogenobenzoyloxy;


R_{2a} is selected from the group consisting of hydrogen, lower alkenyl and lower alkyl;

R_{3a} is selected from the group consisting of hydrogen and lower alkyl;

R_{4a} is selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, fluorine and trifluoromethyl;

R_{5a} is selected from the group consisting of hydrogen, hydroxy, lower alkyl, lower alkoxy, nitro, amino,
 5 lower alkylamino, di(lower alkyl) amino, lower alkanoylamino, lower alkanoyl, lower alkylamino, bis(hydroxy
 lower alkyl)amino, 1-pyrrolidino, 4-methyl-1-piperiziny, 4-morpholiny, cyano, amino lower alkyl, di-lower
 alkyl amino, lower alkyl, trifluoromethyl, halogen, di(lower alkyl)sulfamyl, benzylthio, lower alkylbenzylthio,
 lower alkoxybenzylthio, halogenobenzylthio, benzyloxy, lower alkylbenzyloxy, lower alkoxybenzyloxy,
 10 halogenobenzyloxy, lower alkenyl, lower alkenyloxy, 1-azacyclopropyl, cyclopropyl(lower alkoxy) methoxy,
 and cyclobutyl(lower alkoxy) methoxy;

(b)



wherein:

R_{1b} is hydrogen, lower alkyl or halogenated lower alkyl;

30 R_{2b} is hydrogen or alkyl;

R_{3b}, R_{4b}, R_{5b} and R_{6b} each are hydrogen, lower alkyl, lower alkoxy, nitro, amino, lower alkylamino, lower
 dialkylamino, lower dialkylamino lower alkyl, sulfamyl, lower alkylthio, mercapto, hydroxy, hydroxy lower
 alkyl, lower alkylsulfonyl, halogen, carboxyl, carbo-lower alkoxy, carbamido, halogenoalkyl, cycloalkyl, or
 cycloalkoxy;

35 R_{7b} is lower alkylthio, alkylsulfinyl or alkylsulfonyl;

R_{8b} is hydrogen, halogen, hydroxy, alkoxy or haloalkyl;





40

45


50

55

(c)

 I_{ca}  I_{cb}  I_{cc}  I_{cd}

and

 I_{ce}

wherein:

Ar_c represents a member selected from the group consisting of phenyl, thienyl, 5-methylthienyl, monosubstituted phenyl, disubstituted phenyl and trisubstituted phenyl, each substituent of said substituted phenyls being a member selected from the group consisting of halo, lower alkyl, trifluoromethyl, lower alkoxy, nitro, amino, methylthio and cyano;

Ar_{1c} represents a member selected from the group consisting of phenyl, monosubstituted phenyl, disubstituted phenyl and trisubstituted phenyl, each substituent of said substituted phenyls being a member selected from the group consisting of halo, lower alkyl and lower alkoxy;

R_c represents a member selected from the group consisting of hydrogen and lower alkyl;

R_{1c} represents a member selected from the group consisting of hydrogen, lower alkyl and benzyl;

R_{2c} represents a member selected from the group consisting of




R_{3c} represents a member selected from the group consisting of CO;

R_{4c} represents lower alkyl;


R_{5c} represents lower alkyl; and

R_{6c} represents a member selected from the group consisting of hydrogen and lower alkyl;


(d)




wherein R_d represents a group selected from:



and




(e)

 I_e


wherein R^{1e} and R^{2e} are each radicals of the group consisting of thienyl, furyl, naphthyl, phenyl, and phenyl bearing from one to two substituents of the group consisting of (lower)alkyl, (lower)alkoxy, chloro, bromo, fluoro, di-(lower alkyl)amino, nitro, amino and trifluoromethyl, and wherein R^{3e} is a radical from the group consisting of $-\text{CH}_2\text{-CO}$, $-\text{CH}_2\text{-CH}_2\text{-CO}$, and $-\text{CH}(\text{CH}_3)\text{-CO}$;

(f)

 I_f


wherein Y is hydrogen, chloro, bromo, fluoro, hydroxy, (lower)alkyl, (lower)alkoxy, mercapto, cyano, nitro, amino or (lower)alkylthio.

(g)

 I_g

wherein R_g represents hydrogen or a lower alkyl group having from one to four carbon atoms and R_{1g} represents hydrogen, a lower alkyl group having from one to 4 carbon atoms, a lower alkoxy group having from one to four carbon atoms, chloro, fluoro or bromo, the R_{1g} substitution being at the ortho, meta or para positions of the aroyl group;


(h)



wherein

 R_{1h} represents a hydrogen or halogen atom or a trihalomethyl or lower alkoxy group; R_{2h} represents a hydrogen atom or a lower alkyl group;


15 (i)




wherein

 R_{1i} is a member selected from the group consisting of cycloalkyl with five to seven carbon atoms and lower alkyl substituted cycloalkyl; R_{2i} is a member selected from the group consisting of halogen, nitro, cyano, trifluoromethyl and lower alkylsulfonyl; R_{3i} is hydrogen; X_i is halogen;


35 (j)




wherein R^{1ji} and R^{2ji} are selected from the group consisting of phenyl, halophenyl, lower alkylphenyl, dimethylphenyl, lower alkoxyphenyl, dimethoxyphenyl, lower alkylmercaptophenyl, trifluoromethylphenyl, furyl, thienyl, and naphthyl with the proviso that at least one of R^{1ji} and R^{2ji} is phenyl or substituted phenyl;

wherein R^{1jii} is hydrogen, lower alkyl, phenyl, phenyl monosubstituted with lower alkyl, lower alkoxy or

halogen, naphthyl, benzyl or benzyl wherein the phenyl ring thereof is monosubstituted with lower alkoxy or halogen;




wherein R^{1jiii} is phenyl; R^{2jiii} is phenyl, p-halogenophenyl, p-methylphenyl or furyl; R^{3jiii} is hydrogen, phenyl, or furyl;




25 wherein R^{1jiv} is phenyl, R^{2jiv} is phenyl, p-halogenophenyl, p-methylphenyl, R^{3jiv} is hydrogen or phenyl;

(k)



wherein R_k represents hydrogen or methyl, R_{1k} represents hydrogen, methyl or ethyl, and Y_k represents hydrogen, halogen, or alkyl or alkoxy of 1 through 4 carbon atoms


(l)



55 wherein R_{1l} is a lower alkyl or lower alkyl substituted by cycloalkyl of from three to 12 carbons having from three to seven ring carbons, lower alkenylenyl, lower alkynylenyl, aryl, benzoyl, benzoyl substituted by halogen, lower alkyl, lower alkoxy or heterocycle, such as morpholinyl piperidinyl, furanyl, and thiophenyl; R_{2l} and R_{3l} are the same or different and are lower alkyl, aryl and R_{4l} is hydrogen or lower alkyl

(m)

5

I_m

10

[in which A_m represents a lower straight chain or branched alkylene group;


R_m represents a cyclohexyl or aromatic radical;

15 R'_m represents a halogen atom, trifluoromethyl radical, an alkoxy or alkyl radical containing 1 to 4 carbon atoms or a N,N-dialkylamino radical wherein each of the alkyl groups contain 1 to 4 carbon atoms] as well as alkyl esters thereof (in which the alkyl group in the alcoholic moiety contains 1 to 4 carbon atoms)

(n)

20


25

I_n

30

in which R¹ⁿ is selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower cycloalkyl, phenyl, benzyl and 2-thienyl, R²ⁿ, R³ⁿ, R⁴ⁿ and R⁵ⁿ are the same or different and are each selected from the group consisting of hydrogen and lower alkyl, R^{6an}, R^{6bn}, R^{6cn}, and R^{6dn} are the same or different and selected from the group consisting of hydrogen, lower alkyl, hydroxy, lower alkoxy, benzyloxy, lower alkanoyloxy, nitro, halo, mercapto, lower alkylthio, trifluoromethyl, amino and sulfamoyl, R⁷ⁿ is selected from the group consisting of hydrogen, lower alkyl and lower alkenyl, X is selected from the group consisting of oxy and thio, Y is selected from the group consisting of carbonyl,

40




45

which each of R⁸ⁿ, R⁹ⁿ, R¹⁰ⁿ, R¹¹ⁿ, R¹²ⁿ and R¹³ⁿ is hydrogen or lower alkyl;

(o)

50

I_o

55

wherein R_o is hydrogen or lower alkyl;


(p)




in which R_{1p} is hydrogen or lower alkyl; R_{2p} is hydrogen, alkyl, alkenyl three or four ring-membered cycloalkyl, cycloalkenyl or cycloalkyl-methyl cycloalkenyl-methyl; Ph is (i) 1,3- or 1,4-phenylene, (ii) (lower alkyl)-1,3- or 1,4-phenylene, (iii) (lower alkoxy)-1,3- or 1,4-phenylene, (iv) mono- or di-(halogeno)-1,3- or 1,4-phenylene, (v) (trifluoromethyl)-1,3- or 1,4-phenylene, (vi) (nitro)-1,3- or 1,4-phenylene, (vii) (amino)-1,3- or 1,4-phenylene, (viii) (di-lower alkylamino)-1,3- or 1,4-phenylene; and

$A_p N$ - is monocyclic five to seven ring-membered lower alkyleneamino, piperazino, morpholino, thiamorpholino or N-(lower alkyl, hydroxy lower alkyl, HPh-lower alkyl or HPh) piperazino;

(q)




wherein R_q is hydrogen, halogen, hydroxy, cyano, lower alkyl, hydroxy-lower alkyl, lower alkoxy, acetyl, benzyloxy, lower alkylthio, trifluoromethyl, carboxy, carbo-lower alkoxy, nitro, amino, mono-lower alkylamino, di-lower alkylamino, sulfamoyl, di-lower alkylsulfamoyl or difluoromethylsulfonyl; R_{1q} is halogen, cyano, hydroxy-lower alkyl, lower alkoxy, acetyl, acetamido, benzyloxy, lower alkylthio, trifluoromethyl, hydroxy, carboxy, carbo-lower alkoxy, nitro, amino, mono-lower alkylamino, di-lower alkylamino, sulfamoyl, di-lower alkylsulfamoyl or difluoromethylsulfonyl; or R_q taken together with an adjacent R_{1q} is also lower alkylenedioxy; R_{2q} is




wherein Y_q and X_q , independently, are hydrogen or lower alkyl, and nq is one to seven and R_{3q} is hydrogen, lower alkyl, lower alkoxy-carbonyl-lower alkyl, carboxy-lower alkyl, lower alkanoyl, halo-substituted lower alkanoyl, benzyl, halo-benzyl, benzoyl or halo-benzoyl; and when X_q and Y_q are different, their enantiomers;

(r)

I_r


wherein each of the substituents R^{2r} and R^{3r} is a member of the group consisting of unsubstituted phenyl, naphthyl, thienyl and furyl radicals and phenyl radicals substituted by a substituent selected from the group consisting of halogen, lower alkyl, lower alkoxy, nitro and trifluoromethyl radicals; and wherein R^{1r} is selected from the group consisting of carboxyalkyl- and carboxyalkenyl radicals each containing from two to five carbons;

(s)

I_s

wherein each of X^{1s} and X^{2s} is a hydrogen atom, a halogen atom, an alkyl group having one to four carbon atoms or an alkoxy group having one to four carbon atoms; each of R^{1s} and R^{2s} is a hydrogen atom or an alkyl group having one to four carbon atoms; A^s is carbonyl, methylene or alkylidene having two to four carbon atoms; Y^s is -O-; and ring P_s represents a pyridine or pyridine N-oxide ring;

(t)

I_t

wherein:

Ar_tCO is an aroyl substituent the Ar_t function of which is a member selected from the group consisting of 2-thienyl, 5-lower alkyl-2-thienyl, 5-halo-2-thienyl, 2-naphthyl and 3-pyridyl, said Ar_tCO being in the meta- or para-position relative to the acetyl function;


either of R_t and R_{1t} is hydrogen, the other being a member selected from the group consisting of hydrogen, halo and lower alkyl, provided that, when said R_t is halo or lower alkyl, then said Ar_tCO is in the aforementioned para-position, and when said R_{1t} is halo or lower alkyl, then said Ar_tCO is in the aforementioned meta-position, and further provided that when said R_t or R_{1t} is halo, then said Ar_t is a member selected from the group consisting of 2-thienyl, 5-lower alkyl-2-thienyl and 5-halo-2-thienyl;

either of R_{2t} and R_{3t} is a member selected from the group consisting of hydrogen, allyl and lower alkyl, the other being a member selected from the group consisting of hydrogen and lower alkyl, provided that, when either of said R_{2t} and R_{3t} is allyl, the other is hydrogen, and when either of said R_{2t} and R_{3t} is lower alkyl, the other is a member selected from the group consisting of hydrogen and lower alkyl;

R_{2t} and R_{3t} taken together, is an alkylene bridge attached to the α -carbon of the acetic acid function;

$$\begin{array}{c}
 R_{ti} \\
 | \\
 \text{C}_6\text{H}_3 \\
 | \\
 \text{Ar}_{ti}\text{CO}
 \end{array}
 \begin{array}{c}
 R_{2ti} \\
 | \\
 \text{C} \\
 | \\
 R_{3ti}
 \end{array}
 \begin{array}{c}
 \text{O} \\
 || \\
 \text{C}- \\
 \text{I}_{ti}
 \end{array}$$

R_{2H} and R_{3H} , taken together, is an alkylene bridge attached to the α -carbon of the acetic acid function:


$$\text{Ar}_{\text{tii}}\text{CO}-\text{C}_6\text{H}_3(\text{R}_{\text{1tii}}, \text{R}_{\text{2tii}}, \text{R}_{\text{3tii}})-\text{C}(\text{R}_{\text{2tii}})(\text{R}_{\text{3tii}})\text{C}(=\text{O})\text{I}_{\text{tii}}$$

42

Ar_{tii}Co is an aroyl substituent the Ar_{tii} function of which is a member selected from the group consisting of 2-thienyl, 5-lower alkyl-2-thienyl and 5-halo-2-thienyl, said Ar_{tii}Co being in the meta- or para-position relative to the acetic acid function;


either of R_{tii} and R_{1tii} is hydrogen, the other being a member selected from the group consisting of hydrogen, halo and lower alkyl, provided that, when said R_{tii} is halo or lower alkyl, then said Ar_{tii}Co is in the
5 aforementioned para-position, and when said R_{1tii} is halo or lower alkyl, then said Ar_{tii}Co is in the aforementioned meta-position;

either of R_{2tii} and R_{3tii} is a member selected from the group consisting of hydrogen, allyl and lower alkyl, the other being a member selected from the group consisting of hydrogen and lower alkyl, provided that,
10 when either of said R_{2tii} and R_{3tii} is allyl, the other is hydrogen, and when said R_{2tii} and R_{3tii} is lower alkyl, the other is a member selected from the group consisting of hydrogen and lower alkyl;

R_{2tii} and R_{3tii} together is also an alkylene bridge with α -carbon of the acetic acid function:



wherein n_{tii} is an integer from two to five; and wherein lower alkyl as above employed is a radical having from one to five carbon atoms; or



35 wherein:

Ar_{tiii}Co is an aroyl substituent the Ar_{tiii} function of which is a member selected from the group consisting of 2-thienyl, 5-methyl-2-thienyl and 5-chloro-2-thienyl, said Ar_{tiii}Co being in the meta- or para-position relative to the acetic acid function;

either of R_{tiii} and R_{1tiii} is hydrogen, the other being a member selected from the group consisting of hydrogen, chloro and methyl, provided that, when said R_{tiii} is chloro or methyl, then said Ar_{tiii}Co is in the
40 aforementioned para-position, and when said R_{1tiii} is chloro or methyl, then said Ar_{tiii}Co is in the aforementioned meta-position;


either of R_{2tiii} and R_{3tiii} is a member selected from the group consisting of hydrogen, allyl and lower alkyl, the other being a member selected from the group consisting of hydrogen and lower alkyl, provided that,
45 when either of said R_{2tiii} and R_{3tiii} is allyl, the other is hydrogen, and when either of said R_{2tiii} and R_{3tiii} is lower alkyl, the other is a member selected from the group consisting of hydrogen and lower alkyl;

R_{2tiii} and R_{3tiii} together is also an alkylene bridge with α -carbon of the acetic acid function:



wherein n_{III} is an integer from two to five; and wherein lower alkyl as above employed is a radical having from one to five carbon atoms;

(u)



15 wherein each of R^{1u} and R^{2u} is a hydrogen atom or an alkyl group having one to four carbon atoms; each of X^{1u} and X^{2u} is a hydrogen atom, a halogen atom, an alkyl group having one to four carbon atoms or an alkoxy group having one to four carbon atoms; and Y_u is CO;

(v)




wherein:

35 R_{2v} and R_{3v} each are phenyl or phenyl mono- or di-substituted by at least one member selected from the group consisting of alkyl, alkoxy, alkylmercapto, monoalkylamino, dialkylamino or alkanoylamino wherein the alkyl, alkoxy and alkanoyl each are of up to four carbon atoms, F, Cl, Br, I, CF_3 , OH, methylenedioxy, NH_2 and NO_2 ; A_v is $C_{na}H_{2na}$ wherein na is an integer from one to ten inclusive or



45 and Z_v is O or S;


(w)



where R_{1w}, R_{2w}, R_{3w} and R_{4w} are independently hydrogen, chlorine, fluorine, bromine, lower alkyl, lower alkoxy and R₂ may additionally be trifluoromethyl, and R_{5w} is H or lower alkyl with exclusion of R₁, R₂ and R₃ as hydrogen simultaneously.

20 (x) the acyl residue of (±)-α-methyl-3-phenyl-7-benzofuranepropionic acid;

(y)



where X_y is




35 R_{αy} and R_{βy} are hydrogen or lower alkyl and R_y is halo.

3. A compound according to Claims 1 or 2 which is selected from the group consisting of
- (Z)-6-fluoro-N-hydroxy-N,2-dimethyl-3-[[4-(methylsulfinyl)phenyl]methylene]-3H-indene-1-acetamide;
 - (Z)-6-fluoro-N-methoxy-2-methyl-3-[[4-(methylsulfinyl)phenyl]methylene]-3H-indene-1-acetamide;
 - (Z)-6-fluoro-N-hydroxy-2-methyl-3-[[4-(methylsulfinyl)phenyl]methylene]-3H-indene-1-acetamide;
 - 40 (Z)-6-fluoro-N-hydroxy-N-(1-methylethyl)-2-(methyl-3-[[4-(methylsulfinyl)phenyl]methylene]-3H-indene-1-acetamide;
 - 1-(4-chlorobenzoyl-N-hydroxy-5-methoxy-N,2-dimethyl-1H-indole-3-acetamide;
 - 1-(4-chlorobenzoyl-N,5-dimethoxy-2-methyl-1H-indole-3-acetamide;
 - 1-(4-chlorobenzoyl-N-hydroxy-5-methoxy-2-methyl-N-(1-methylethyl)-1H-indole-3-acetamide;
 - 45 1-(4-chlorobenzoyl-N-cyclohexyl-N-hydroxy-5-methoxy-2-methyl-1H-indole-3-acetamide;
 - 1-(4-chlorobenzoyl-N,5-dimethoxy-N,2-dimethyl-1H-indole-3-acetamide;
 - 1-(4-chlorobenzoyl-N-hydroxy-5-methoxy-2-methyl-N-phenyl-1H-indole-3-acetamide;
 - 1-(4-chlorobenzoyl-N-benzyl-N-hydroxy-5-methoxy-2-methyl-1H-indole-3-acetamide;
 - N-methoxy-2-[(2,6-dichlorophenyl)amino]benzeneacetamide or
 - 50 N-hydroxy-N-methyl-2-[(2,6-dichlorophenyl)amino]benzeneacetamide.

4. A compound according to Claims 1 to 2

- wherein I_c is 5-(4-methylbenzoyl)-1-methylpyrrolo-2-acetyl; or
- wherein RCO is I_a wherein R^{1a} is phenyl, R^{3a} is -CH₂CO- and R^{2a} is p-chlorophenyl; or
- wherein RCO is I_f wherein Y is chloro; or
- 55 wherein RCO is I_g wherein R_g and R_{1g} are hydrogen; or
- wherein RCO is I_h wherein R_{1h} and R_{2h} are hydrogen; or
- wherein RCO is I_i wherein R_{1i} is cyclohexyl, R_{3i} is H and R_{2i} and X_i are chloro; or
- wherein RCO is I_j wherein R^{1j} is phenyl, and R^{2j} is para-chlorophenyl; or


wherein RCO is I_k wherein Y_k and R_{1k} are hydrogen and R_k is methyl; or
 wherein RCO is I_l wherein R_{2l} and R_{3l} are CH_3 , R_{4l} is hydrogen and R_{1l} is p-chlorophenyl; or
 wherein RCO is I_m wherein R_m is methoxy, A_m is $-CH_2-$, and R_m is p-chlorophenyl; or
 wherein RCO is I_n wherein R^{1n} and R^{6n} are ethyl, Y is acetyl, and R^{2n} , R^{3n} , R^{4n} , R^{5n} , R^{6an} , R^{6bn} , R^{6cn} and R^{7n}
 5 are hydrogen; or
 wherein RCO is I_o wherein R_o is CH_3 ; or
 wherein RCO is I_p wherein one of R_{1p} is hydrogen and one of R_{2p} is methyl, Ph is -1,4-(3-chlorophenylene)
 and A_pN- is




15 or
 wherein RCO is I_q wherein R_{2q} is 2-acetyl, R_q is 6-chloro, and R_{1q} and R_{3q} are hydrogen; or
 wherein RCO is I_r wherein R^{1r} is 3-propionyl and R^{2r} and R^{3r} are phenyl; or
 wherein RCO is 2-(5H-[1]benzopyrano[2,3b]pyridin-7-yl) propionyl; or
 wherein RCO is I_t wherein A_t is 2-thienyl, R_t , R_{1t} and R_{2t} are hydrogen and R_{3t} is methyl; or
 20 wherein RCO is I_u wherein X^{1u} , X^{2u} and R^{1u} are hydrogen and Y^{2u} is methyl; or
 wherein RCO is I_v wherein Z_v is oxygen, A_v is



and R_{2v} and R_{3v} are each p-chlorophenyl; or
 wherein RCO is I_y wherein R_y is m-chloro and $R_{\alpha y}$ and $R_{\beta y}$ are hydrogen
 30 5. A pharmaceutical composition comprising an effective amount of a compound of the formula I of
 Claim 1 together with a pharmaceutically acceptable carrier.
 6. A method of use of a compound according to Claim 1-4 for the manufacturing of pharmaceuticals for
 treating a condition advantageously affected by inhibition of 5-lipoxygenase, cyclooxygenase, or both.
 7. A process for the preparation of a compound of the formula I of Claim 1 or 2 wherein a compound of
 35 the formula
 $RCOOH$
 wherein RCO is as defined above in step (1) treated with a compound selected from the group consisting of
 (i) bromo or chloro producing agent in the presence of a base; or
 (ii) a coupling agent;
 40 and then the product of step (1) is further treated with a compound of the formula



wherein R_1 and R_2 are as defined above to obtain the compound of the formula I.
 50 8. A process for the preparation of a compound of the formula I of Claim 2 wherein a compound of the
 formula
 $RCOOH$
 wherein RCO is as defined above is (1) treated with a coupling agent and then (2) reacted with a compound
 55 of the formula



5

wherein R_1 and R_2 are as defined above to obtain the compound of the formula I.

9. A process according to Claims 7 or 8 for the preparation of a compound of formula I wherein RCO is I_b wherein R_{7b} is alkylsulfinyl or wherein RCO is I_w , or wherein RCO is I_b wherein R_{7b} is alkylsulfinyl; or wherein RCO is I_w .


10

Claims for the following Contracting States: ES, GR

15

1. A process for the preparation of a compound of the formula

20



I

25 and pharmaceutically acceptable acid addition or base salts thereof wherein


i) RCO is an acyl residue of selected NSAIDs;

ii) R_1 is hydrogen, lower alkyl or acyl;

30 iii) R_2 is H, lower alkyl, cycloalkyl of from three to twelve carbon atoms of which from three to seven carbons are ring carbons, aryl, arylalkyl or heteroaryl with the overall proviso that when the acyl residue is of the formula

30

35




40

wherein R_4 is hydrogen or lower alkyl, then R_3 is hydrogen, phenylmethyl, allyl, vinyl, isopropenyl, benzoyl, chlorobenzoyl, methoxybenzoyl, and thiomethylbenzoyl; and when the acyl residue is of the formula

45

50


55



wherein R_5 , R_6 , R_7 and R_8 are independently H, chlorine, fluorine, or bromine, lower alkyl, lower alkoxy, or trifluoromethyl but when R_4 is hydrogen then are not all hydrogen; and R_4 is otherwise independently as defined above; then R_1 and R_2 cannot both be hydrogen, comprising a compound of the formula $RCOOH$

wherein RCO is as defined above in step (1) treated with a compound selected from the group consisting of (i) bromo or chloro producing agent in the presence of a base; or (ii) a coupling agent;


and then the product of step (1) is further treated with a compound of the formula



wherein R_1 are as defined above or an acid addition salt thereof, to obtain the compound of the formula I.

2. A process according to claim 1 for the preparation of a compound of formula I wherein RCO is selected from the groups consisting of

(a)



in which


R_{1a} is selected from the group consisting of benzene, naphthalene, biphenyl and substituted benzene, naphthalene and biphenyl radicals in which said substituent is selected from the group consisting of halogen, lower alkyl, lower alkylthio, lower alkoxy, trifluoromethyl, phenoxy, lower alkyl phenoxy, lower alkoxy phenoxy, halogenophenoxy, trifluoroacetyl, difluoroacetyl, monofluoroacetyl, di-lower alkyl sulfamyl, lower alkanoyl, di-lower alkyl carboxamido, cyano, carb-lower alkoxy, trifluoromethylthio, lower alkyl sulfinyl, lower alkylsulfonyl, benzylthio, lower alkylbenzylthio, lower alkoxybenzylthio, halogenobenzylthio, mercapto, nitro, amino, di-(lower alkyl)amino, lower alkylamino, lower alkanoylamino, hydroxy, lower alkanoyloxy, trifluoroacetoxyl, difluoroacetoxyl, monofluoroacetoxyl, benzyloxy, lower alkylbenzyloxy, lower alkoxybenzyloxy, and halogenobenzyloxy;

R_{2a} is selected from the group consisting of hydrogen, lower alkenyl and lower alkyl;

R_{3a} is selected from the group consisting of hydrogen and lower alkyl;

R_{4a} is selected from the group consisting of hydrogen, lower alkyl, lower alkoxy, fluorine and trifluoromethyl;
 R_{5a} is selected from the group consisting of hydrogen, hydroxy, lower alkyl, lower alkoxy, nitro, amino,
 lower alkylamino, di(lower alkyl) amino, lower alkanoylamino, lower alkanoyl, lower alkylamino, bis(hydroxy
 lower alkyl)amino, 1-pyrrolidino, 4-methyl-1-piperiziny, 4-morpholiny, cyano, amino lower alkyl, di-lower
 5 alkyl amino, lower alkyl, trifluoromethyl, halogen, di(lower alkyl)sulfamyl, benzylthio, lower alkylbenzylthio,
 lower alkoxybenzylthio, halogenobenzylthio, benzyloxy, lower alkylbenzyloxy, lower alkoxybenzyloxy,
 halogenobenzyloxy, lower alkenyl, lower alkenyloxy, 1-azacyclopropyl, cyclopropyl(lower alkoxy) methoxy,
 and cyclobutyl(lower alkoxy) methoxy;

10 (b)



25

wherein

R_{1b} is hydrogen, lower alkyl or halogenated lower alkyl;

R_{2b} is hydrogen or alkyl;

R_{3b} , R_{4b} , R_{5b} and R_{6b} each are hydrogen, lower alkyl, lower alkoxy, nitro, amino, lower alkylamino, lower
 30 dialkylamino, lower dialkylamino lower alkyl, sulfamyl, lower alkylthio, mercapto, hydroxy, hydroxy lower
 alkyl, lower alkylsulfonyl, halogen, carboxyl, carbo-lower alkoxy, carbamido, halogenoalkyl, cycloalkyl, or
 cycloalkoxy;

R_{7b} is lower alkylthio, alkylsulfinyl or alkylsulfonyl;

35 R_{8b} is hydrogen, halogen, hydroxy, alkoxy or haloalkyl;




40

45


50

55

(c)

 I_{ca}  I_{cb}  I_{cc}  I_{cd}

and

 I_{ce}

wherein:

Ar_c represents a member selected from the group consisting of phenyl, thienyl, 5-methylthienyl, monosubstituted phenyl, disubstituted phenyl and trisubstituted phenyl, each substituent of said substituted phenyls being a member selected from the group consisting of halo, lower alkyl, trifluoromethyl, lower alkoxy, nitro, amino, methylthio and cyano;

Ar_{1c} represents a member selected from the group consisting of phenyl, monosubstituted phenyl, disubstituted phenyl and trisubstituted phenyl, each substituent of said substituted phenyls being a member selected from the group consisting of halo, lower alkyl and lower alkoxy;

R_c represents a member selected from the group consisting of hydrogen and lower alkyl;

R_{1c} represents a member selected from the group consisting of hydrogen, lower alkyl and benzyl;

R_{2c} represents a member selected from the group consisting of




R_{3c} represents a member selected from the group consisting of CO;

R_{4c} represents lower alkyl;


R_{5c} represents lower alkyl; and

R_{6c} represents a member selected from the group consisting of hydrogen and lower alkyl;


(d)




wherein R_d represents a group selected from:



and




(e)

 I_e


wherein R^{1e} and R^{2e} are each radicals of the group consisting of thienyl, furyl, naphthyl, phenyl, and phenyl bearing from one to two substituents of the group consisting of (lower)alkyl, (lower)alkoxy, chloro, bromo, fluoro, di-(lower alkyl)amino, nitro, amino and trifluoromethyl, and wherein R^{3e} is a radical from the group consisting of $-\text{CH}_2-\text{CO}$, $-\text{CH}_2-\text{CH}_2-\text{CO}$, and $-\text{CH}(\text{CH}_3)-\text{CO}$;

(f)

 I_f


wherein Y is hydrogen, chloro, bromo, fluoro, hydroxy, (lower)alkyl, (lower)alkoxy, mercapto, cyano, nitro, amino or (lower)alkylthio;

(g)

 I_g

wherein R_g represents hydrogen or a lower alkyl group having from one to four carbon atoms and R_{1g} represents hydrogen, a lower alkyl group having from one to 4 carbon atoms, a lower alkoxy group having from one to four carbon atoms, chloro, fluoro or bromo, the R_{1g} substitution being at the ortho, meta or para positions of the aroyl group;


(h)

 I_h

wherein

 R_{1h} represents a hydrogen or halogen atom or a trihalomethyl or lower alkoxy group; R_{2h} represents a hydrogen atom or a lower alkyl group;


(i)

 I_i


wherein

 R_{1i} is a member selected from the group consisting of cycloalkyl with five to seven carbon atoms and lower alkyl substituted cycloalkyl; R_{2i} is a member selected from the group consisting of halogen, nitro, cyano, trifluoromethyl and lower alkylsulfonyl; R_{3i} is hydrogen; X_i is halogen;

(j)


 I_{ji}

wherein R^{1ji} and R^{2ji} are selected from the group consisting of phenyl, halophenyl, lower alkylphenyl, dimethylphenyl, lower alkoxyphenyl, dimethoxyphenyl, lower alkylmercaptophenyl, trifluoromethylphenyl, furyl, thienyl, and naphthyl with the proviso that at least one of R^{1ji} and R^{2ji} is phenyl or substituted phenyl;


 I_{jii}

wherein R^{1jii} is hydrogen, lower alkyl, phenyl, phenyl monosubstituted with lower alkyl, lower alkoxy or halogen, naphthyl, benzyl or benzyl wherein the phenyl ring thereof is monosubstituted with lower alkoxy or

halogen;




wherein R^{1jiii} is phenyl; R^{2jiii} is phenyl, p-halogenophenyl, p-methylphenyl or furyl; R^{3jiii} is hydrogen, phenyl, or furyl;




25 wherein R^{1jiv} is phenyl, R^{2jiv} is phenyl, p-halogenophenyl, p-methylphenyl, R^{3jiv} is hydrogen or phenyl;

(k)




40 wherein R_k represents hydrogen or methyl, R_{1k} represents hydrogen, methyl or ethyl, and Y_k represents hydrogen, halogen, or alkyl or alkoxy of 1 through 4 carbon atoms;

(l)



55 wherein R_{11} is a lower alkyl or lower alkyl substituted by cycloalkyl of from three to 12 carbons having from three to seven ring carbons, lower alkenylenyl, lower alkynylenyl, aryl, benzoyl, benzoyl substituted by halogen, lower alkyl, lower alkoxy or heterocycle, such as morpholinyl piperidinyl, furanyl, and thiophenyl; R_{2a} and R_{3a} are the same or different and are lower alkyl, aryl and R_{4a} is hydrogen or lower alkyl;

(m)




[in which A_m represents a lower straight chain or branched alkylene group;


R_m represents a cyclohexyl or aromatic radical;

15 R'_m represents a halogen atom, trifluoromethyl radical, an alkoxy or alkyl radical containing 1 to 4 carbon atoms or a N,N-dialkylamino radical wherein each of the alkyl groups contain 1 to 4 carbon atoms] as well as alkyl esters thereof (in which the alkyl group in the alcoholic moiety contains 1 to 4 carbon atoms);

(n)




30 in which R^{1n} is selected from the group consisting of lower alkyl, lower alkenyl, lower alkynyl, lower cycloalkyl, phenyl, benzyl and 2-thienyl, R^{2n} , R^{3n} , R^{4n} and R^{5n} are the same or different and are each selected from the group consisting of hydrogen and lower alkyl, R^{6an} , R^{6bn} , R^{6cn} , and R^{6dn} are the same or different and selected from the group consisting of hydrogen, lower alkyl, hydroxy, lower alkoxy, benzyloxy, lower alkanoyloxy, nitro, halo, mercapto, lower alkylthio, trifluoromethyl, amino and sulfamoyl, R^{7n} is selected from the group consisting of hydrogen, lower alkyl and lower alkenyl, X is selected from the group consisting of oxy and thio, Y is selected from the group consisting of carbonyl,



which each of R^{8n} , R^{9n} , R^{10n} , R^{11n} , R^{12n} and R^{13n} is hydrogen or lower alkyl;

(o)



wherein R_p is hydrogen or lower alkyl;

(p)




in which R_{1p} is hydrogen or lower alkyl; R_{2p} is hydrogen, alkyl, alkenyl three or four ring-membered cycloalkyl, cycloalkenyl or cycloalkyl-methyl cycloalkenyl-methyl; Ph is (i) 1,3- or 1,4-phenylene, (ii) (lower alkyl)-1,3- or 1,4-phenylene, (iii) (lower alkoxy)-1,3- or 1,4-phenylene, (iv) mono- or di-(halogeno)-1,3- or 1,4-phenylene, (v) (trifluoromethyl)-1,3- or 1,4-phenylene, (vi) (nitro)-1,3- or 1,4-phenylene, (vii) (amino)-1,3- or 1,4-phenylene, (viii) di-lower alkylamino)-1,3- or 1,4-phenylene; and


$A_p N-$

is monocyclic five to seven ring-membered lower alkyleneamino, piperazino, morpholino, thiamorpholino or N-(lower alkyl, hydroxy lower alkyl, HPh-lower alkyl or HPh) piperazino;

(q)




wherein R_q is hydrogen, halogen, hydroxy, cyano, lower alkyl, hydroxy-lower alkyl, lower alkoxy, acetyl, benzyloxy, lower alkylthio, trifluoromethyl, carboxy, carbo-lower alkoxy, nitro, amino, mono-lower alkylamino, di-lower alkylamino, sulfamoyl, di-lower alkylsulfamoyl or difluoromethylsulfonyl; R_{1q} is halogen, cyano, hydroxy-lower alkyl, lower alkoxy, acetyl, acetamido, benzyloxy, lower alkylthio, trifluoromethyl, hydroxy, carboxy, carbo-lower alkoxy, nitro, amino, mono-lower alkylamino, di-lower alkylamino, sulfamoyl, di-lower alkylsulfamoyl or difluoromethylsulfonyl; or R_q taken together with an adjacent R_{1q} is also lower alkylendioxy; R_{2q} is




wherein Y_q and X_q , independently, are hydrogen or lower alkyl, and n_q is one to seven and R_{3q} is hydrogen, lower alkyl, lower alkoxy-carbonyl-lower alkyl, carboxy-lower alkyl, lower alkanoyl, halo-substituted lower alkanoyl, benzyl, halo-benzyl, benzoyl or halo-benzoyl; and when X_q and Y_q are different, their enantiomers;

(r)

I_r


wherein each of the substituents R^{2r} and R^{3r} is a member of the group consisting of unsubstituted phenyl, naphthyl, thienyl and furyl radicals and phenyl radicals substituted by a substituent selected from the group consisting of halogen, lower alkyl, lower alkoxy, nitro and trifluoromethyl radicals; and wherein R^{1r} is selected from the group consisting of carboxyalkyl- and carboxyalkenyl radicals each containing from two to five carbons;

(s)

I_s

wherein each of X^{1s} and X^{2s} is a hydrogen atom, a halogen atom, an alkyl group having one to four carbon atoms or an alkoxy group having one to four carbon atoms; each of R^{1s} and R^{2s} is a hydrogen atom or an alkyl group having one to four carbon atoms; A^s is carbonyl, methylene or alkylidene having two to four carbon atoms; Y_s is -O-; and ring P_s represents a pyridine or pyridine N-oxide ring;

(t)

I_t


wherein:

Ar_tCO is an aroyl substituent the Ar_t function of which is a member selected from the group consisting of 2-thienyl, 5-lower alkyl-2-thienyl, 5-halo-2-thienyl, 2-naphthyl and 3-pyridyl, said Ar_tCO being in the meta- or para-position relative to the acetyl function;


either of R_t and R_{1t} is hydrogen, the other being a member selected from the group consisting of hydrogen, halo and lower alkyl, provided that, when said R_t is halo or lower alkyl, then said Ar_tCO is in the aforementioned para-position, and when said R_{1t} is halo or lower alkyl, then said Ar_tCO is in the aforementioned meta-position, and further provided that when said R_t or R_{1t} is halo, then said Ar_t is a member selected from the group consisting of 2-thienyl, 5-lower alkyl-2-thienyl and 5-halo-2-thienyl;

either of R_{2t} and R_{3t} is a member selected from the group consisting of hydrogen, allyl and lower alkyl, the other being a member selected from the group consisting of hydrogen and lower alkyl, provided that, when either of said R_{2t} and R_{3t} is allyl, the other is hydrogen, and when either of said R_{2t} and R_{3t} is lower alkyl, the other is a member selected from the group consisting of hydrogen and lower alkyl;

R_{2t} and R_{3t} taken together, is an alkylene bridge attached to the α-carbon of the acetic acid function:



10 wherein n_t is an integer from two to five; and wherein lower alkyl as above employed is a radical having from one to five carbon atoms;




wherein

25 $Ar_{ti}CO$ is an aroyl substituent the Ar_{ti} function of which is a member selected from the group consisting of 2-thienyl, 5-methyl-2-thienyl, 5-chloro-2-thienyl, 2-naphthyl and 3-pyridyl, said $Ar_{ti}CO$ being in the meta- or para-position relative to the acetic acid function;


30 either of R_{ti} and R_{1ti} is hydrogen, the other being a member selected from the group consisting of hydrogen, chloro and methyl, provided that, when said R_{ti} is chloro or methyl, then said $Ar_{ti}CO$ is in the aforementioned para-position, and when said R_{1ti} is chloro or methyl, then said $Ar_{ti}CO$ is in the aforementioned meta-position, and further provided that, when said R_{ti} and R_{1ti} is chloro, then said Ar_{ti} is a member selected from the group consisting of 2-thienyl, 5-methyl-2-thienyl and 5-chloro-2-thienyl;

35 either of R_{2ti} and R_{3ti} is a member selected from the group consisting of hydrogen, allyl and lower alkyl, the other being a member selected from the group consisting of hydrogen and lower alkyl, provided that, when either of said R_{2ti} and R_{3ti} is allyl, the other is hydrogen, and when either of said R_{2ti} and R_{3ti} is lower alkyl, the other is a member selected from the group consisting of hydrogen and lower alkyl;

R_{2ti} and R_{3ti} , taken together, is an alkylene bridge attached to the α -carbon of the acetic acid function:



wherein n_{ti} is an integer from two to five; and wherein lower alkyl as above employed is a radical having from one to five carbons;




wherein:

15 $Ar_{tii}CO$ is an aroyl substituent the Ar_{tii} function of which is a member selected from the group consisting of 2-thienyl, 5-lower alkyl-2-thienyl and 5-halo-2-thienyl, said $Ar_{tii}CO$ being in the meta- or para-position relative to the acetic acid function;


either of R_{tii} and R_{ltii} is hydrogen, the other being a member selected from the group consisting of hydrogen, halo and lower alkyl, provided that, when said R_{tii} is halo or lower alkyl, then said $Ar_{tii}CO$ is in the aforementioned para-position, and when said R_{ltii} is halo or lower alkyl, then said $Ar_{tii}CO$ is in the

20 aforementioned meta-position;
either of R_{2tii} and R_{3tii} is a member selected from the group consisting of hydrogen, allyl and lower alkyl, the other being a member selected from the group consisting of hydrogen and lower alkyl, provided that, when either of said R_{2tii} and R_{3tii} is allyl, the other is hydrogen, and when said R_{2tii} and R_{3tii} is lower alkyl, the other is a member selected from the group consisting of hydrogen and lower alkyl;

25 R_{2tii} and R_{3tii} together is also an alkylene bridge with α -carbon of the acetic acid function:



35 wherein n_{tii} is an integer from two to five; and wherein lower alkyl as above employed is a radical having from one to five carbon atoms; or



wherein:

50 $Ar_{tiii}CO$ is an aroyl substituent the Ar_{tiii} function of which is a member selected from the group consisting of 2-thienyl, 5-methyl-2-thienyl and 5-chloro-2-thienyl, said $Ar_{tiii}CO$ being in the meta- or para-position relative to the acetic acid function;


either of R_{tiii} and R_{ltiii} is hydrogen, the other being a member selected from the group consisting of hydrogen, chloro and methyl, provided that, when said R_{tiii} is chloro or methyl, then said $Ar_{tiii}CO$ is in the aforementioned para-position, and when said R_{ltiii} is chloro or methyl, then said $Ar_{tiii}CO$ is in the

55 aforementioned meta-position;
either of R_{2tiii} and R_{3tiii} is a member selected from the group consisting of hydrogen, allyl and lower alkyl, the other being a member selected from the group consisting of hydrogen and lower alkyl, provided that,

when either of said $R_{2\text{tiii}}$ and $R_{3\text{tiii}}$ is allyl, the other is hydrogen, and when either of said $R_{2\text{tiii}}$ and $R_{3\text{tiii}}$ is lower alkyl, the other is a member selected from the group consisting of hydrogen and lower alkyl; $R_{2\text{tiii}}$ and $R_{3\text{tiii}}$ together is also an alkylene bridge with α -carbon of the acetic acid function:

5

10




wherein n_{tiii} is an integer from two to five; and wherein lower alkyl as above employed is a radical having from one to five carbon atoms;

15

(u)

20



25


wherein each of R^{1u} and R^{2u} is a hydrogen atom or an alkyl group having one to four carbon atoms; each of X^{1u} and X^{2u} is a hydrogen atom, a halogen atoms, an alkyl group having one to four carbon atoms or an alkoxy group having one to four carbon atoms; and Y_u is CO;

30

(v)

35

40

 I_v

45 wherein:

R_{2v} and R_{3v} each are phenyl or phenyl mono- or di-substituted by at least one member selected from the group consisting of alkyl, alkoxy, alkylmercapto, monoalkylamino, dialkylamino or alkanoylamino wherein the alkyl, alkoxy and alkanoyl each are of up to four carbon atoms, F, Cl, Br, I, CF_3 , OH, methylenedioxy, NH_2 and NO_2 ;


50 A_v is $C_{na}H_{2na}$ wherein na is an integer from one to ten inclusive or

55



and Z_v is O or S;


(w)



where R_{1w} , R_{2w} , R_{3w} and R_{4w} are independently hydrogen, chlorine, fluorine, bromine, lower alkyl, lower alkoxy and R_2 may additionally be trifluoromethyl, and R_{5w} is H or lower alkyl with exclusion of R_1 , R_2 and R_3 as hydrogen simultaneously;

20 (x) the acyl residue of (\pm)- α -methyl-3-phenyl-7-benzofuranepropionic acid;

(y)



where X_y is



35 $R_{\alpha y}$ and $R_{\beta y}$ are hydrogen or lower alkyl and R_y is halo.

3. A process according to claim 1 for the preparation of a compound of formula I which is selected from the group consisting of


- (Z)-6-fluoro-N-hydroxy-N,2-dimethyl-3-[[4-(methylsulfinyl)phenyl]methylene]-3H-indene-1-acetamide;
 (Z)-6-fluoro-N-methoxy-2-methyl-3-[[4-(methylsulfinyl)phenyl]methylene]-3H-indene-1-acetamide;
 40 (Z)-6-fluoro-N-hydroxy-2-methyl-3-[[4-(methylsulfinyl)phenyl]methylene]-3H-indene-1-acetamide;
 (Z)-6-fluoro-N-hydroxy-N-(1-methylethyl)-2-methyl-3-[[4-(methylsulfinyl)phenyl]methylene]-3H-indene-1-acetamide;
 1-(4-chlorobenzoyl)-N-hydroxy-5-methoxy-N,2-dimethyl-1H-indole-3-acetamide;
 1-(4-chlorobenzoyl)-N,5-dimethoxy-2-methyl-1H-indole-3-acetamide;
 45 1-(4-chlorobenzoyl)-N-hydroxy-5-methoxy-2-methyl-N-(1-methylethyl)-1H-indole-3-acetamide;
 1-(4-chlorobenzoyl)-N-cyclohexyl-N-hydroxy-5-methoxy-2-methyl-1H-indole-3-acetamide;
 1-(4-chlorobenzoyl)-N,5-dimethoxy-N,2-dimethyl-1H-indole-3-acetamide;
 1-(4-chlorobenzoyl)-N-hydroxy-5-methoxy-2-methyl-N-phenyl-1H-indole-3-acetamide;
 1-(4-chlorobenzoyl)-N-benzyl-N-hydroxy-5-methoxy-2-methyl-1H-indole-3-acetamide;
 50 N-methoxy-2-[(2,6-dichlorophenyl)amino]benzeneacetamide or N-hydroxy-N-methyl-2-[(2,6-dichlorophenyl)amino] benzeneacetamide.

4. A process for the preparation of a compound of formula I according to claim 2

- wherein I_c is 5-(4-methylbenzoyl)-1-methylpyrrolo-2-acetyl; or
 wherein RCO is I_e wherein R^{1e} is phenyl, R^{3e} is $-\text{CH}_2\text{CO}-$ and R^{2e} is p-chlorophenyl; or
 wherein RCO is I_f wherein Y is chloro; or
 55 wherein RCO is I_g wherein R_g and R_{1g} are hydrogen; or
 wherein RCO is I_h wherein R_{1h} and R_{2h} are hydrogen; or
 wherein RCO is I_i wherein R_{1i} is cyclohexyl, R_{3i} is H and R_{2i} and X_i are chloro; or

wherein RCO is I_{ij} wherein R^{1j} is phenyl, and R^{2j} is para-chlorophenyl; or
 wherein RCO is I_k wherein Y_k and R_{1k} are hydrogen and R_k is methyl; or
 wherein RCO is I_l wherein R_{2l} and R_{3l} are CH_3 , R_{4l} is hydrogen and R_{1l} is p-chlorophenyl; or
 wherein RCO is I_m wherein R_m is methoxy, A_m is $-CH_2-$, and R_m is p-chlorophenyl; or
 5 wherein RCO is I_n wherein R^{1n} and R^{6dn} are ethyl, Y is acetyl, and R^{2n} , R^{3n} , R^{4n} , R^{5n} , R^{6an} , R^{6bn} , R^{6cn} and R^{7n} are hydrogen; or
 wherein RCO is I_o wherein R_o is CH_3 ; or
 wherein RCO is I_p wherein one of R_{1p} is hydrogen and one of R_{2p} is methyl, Ph is -1,4-(3-chlorophenylene)
 and A_pN- is or

10



15

wherein RCO is I_q wherein R_{2q} is 2-acetyl, R_q is 6-chloro, and R_{1q} and R_{3q} are hydrogen; or
 wherein RCO is I_r wherein R^{1r} is 3-propionyl and R^{2r} and R^{3r} are phenyl; or
 wherein RCO is 2-(5H-[1]benzopyrano[2,3b]pyridin-7-yl) propionyl; or
 20 wherein RCO is I_t wherein A_t is 2-thienyl, R_t , R_{1t} and R_{2t} are hydrogen and R_{3t} is methyl; or
 wherein RCO is I_u wherein X^{1u} , X^{2u} and R^{1u} are hydrogen and Y^{2u} is methyl; or
 wherein RCO is I_v wherein Z_v is oxygen, A_v is

25



and R_{2v} and R_{3v} are each p-chlorophenyl; or
 30 wherein RCO is I_y wherein R_y is m-chloro and $R_{\alpha y}$ and $R_{\beta y}$ are hydrogen.


5. A method of use of a compound of formula I according to claim 1 for the manufacturing of pharmaceuticals for treating a condition advantageously affected by inhibition of 5-lipoxygenase, cyclooxygenase, or both.

6. A process according to claim 1 for the preparation of a compound of the formula I wherein a
 35 compound of the formula

RCOOH

wherein RCO is as defined above is (1) treated with a coupling agent and then (2) reacted with a compound of the formula

40



45

wherein R_1 and R_2 are as defined above to obtain the compound of the formula I.

7. A process of the Claim 1 wherein RCO is I_b wherein R_{7b} is alkylsulfinyl or wherein RCO is I_w .

8. A process of the Claim 1 wherein RCO is I_b wherein R_{7b} is alkylsulfinyl.

50 9. A process of the Claim 1 wherein RCO is I_w .

55



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number:

0 342 682 A3

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: **89108976.5**

(51) Int. Cl.⁵: **C07D 209/28, C07C 83/10,
A61K 31/405, A61K 31/16**

(22) Date of filing: **18.05.89**

(30) Priority: **19.05.88 US 196584**

(43) Date of publication of application:
23.11.89 Bulletin 89/47

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(88) Date of deferred publication of the search report:
11.12.91 Bulletin 91/50


(71) Applicant: **WARNER-LAMBERT COMPANY**
201 Tabor Road
Morris Plains New Jersey 07950(US)

(72) Inventor: **Cetenko, Wiaczeslaw Antin**
3565 Green Brier
Ann Arbor Michigan 48105(US)
Inventor: **Connor, David Thomas**
2453 Antietam
Ann Arbor Michigan 48105(US)
Inventor: **Flynn, Daniel Lee**
17 North Bristol Court
Mundelein Illinois 60060(US)
Inventor: **Sircar, Jagadish Chandra**
3615 Charter Place
Ann Arbor, Michigan 48105(US)

(74) Representative: **Mansmann, Ivo**
c/o Gödecke AG - Patentabteilung Postfach
569 Mooswaldallee 1-9
W-7800 Freiburg(DE)

(54) **Novel hydroxamate derivatives of selected nonsteroidal anti-inflammatory acyl residues having cyclooxygenase and 5-lipoxygenase inhibition.**

(57) A compound of the formula



and pharmaceutically acceptable acid addition or base salts thereof wherein

i) RCO is an acyl residue of selected NSAIDS;

ii) R₁ is hydrogen, lower alkyl or acyl;

iii) R₂ is H, lower alkyl, cycloalkyl of from three to twelve carbon atoms of which from three to seven carbons are ring carbons, aryl, arylalkyl or heteroaryl having cyclooxygenase or 5-lipoxygenase inhibitory activity.

EP 0 342 682 A3



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

which under Rule 45 of the European Patent Convention
shall be considered, for the purposes of subsequent
proceedings, as the European search report

Application Number

EP 89 10 8976

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
X	ARZNEIMITTEL FORSCHUNG DRUG RESEARCH, vol. 30(II), no. 8a, August 1980, pages 1314-1325, Aulendorf, DE; K.-H. BOLTZE et al.: "Chemische Struktur und antiphlogistische Wirkung in der Reihe der substituierten Indol-3-essigsäuren" * Page 1317, table 2, compound no. 40; page 1319, column 1, 2nd paragraph *	1-3,6	C 07 D 209/28 C 07 C 83/10 A 61 K 31/405 A 61 K 31/16
X	DE-A-2 008 332 (ISTITUTO BIOLOGICO CHEMIOTERAPICO ABC S.p.A.) * Pages 15-16, claims 1-4; pages 5-6, example 3; page 14, 1st paragraph *	1,2,6,7	
X	FR-A-1 579 495 (R. ARIES) * Page 2, column 1, examples 2-4; page 1, column 2, 7th paragraph *	1,2,6	
A	EP-A-0 039 051 (MERCK & CO. INC.) * The whole document; especially pages 17-19, examples 3-5 *	1-4,6	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			C 07 D 209/00 C 07 C 83/00 C 07 C 147/00
INCOMPLETE SEARCH			
<p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art of some of the claims</p> <p>Claims searched completely: 3, 5-9 Claims searched incompletely: 1, 2 Claims not searched: 4</p> <p>Reason for the limitation of the search:</p> <p>See sheet -c-</p>			
Place of search		Date of completion of the search	Examiner
THE HAGUE		23-08-1991	FINK D.
CATEGORY OF CITED DOCUMENTS			
<p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

EP 89 10 8976

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
D,A	<p>JOURNAL OF MEDICINAL CHEMISTRY, vol. 30, no. 3, March 1987, pages 574-580, American Chemical Society, Columbus, Ohio, US; J.B. SUMMERS et al.: "Hydroxamic acid inhibitors of 5-lipoxygenase"</p> <p>* The whole article; especially page 575, column 1, last paragraph - column 2, 1st paragraph *</p> <p>-----</p>	1-4,6	
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)



EP 89 10 8976 -C-

Claims 1 and 2 are much too broad with respect to the definition of the NSAIDS-residue. Hence, for economic reasons (see Guidelines for Examination in the EPO, Part B, III 2.1 and III 2.3) claims 1 and 2 cannot be searched completely. Moreover the said claims are only partly supported by the descriptive part of the application (Art. 84 EPC). So the search has been restricted to those classes of compounds for which real examples exist (claim 2(a), 2(b) and 2(w)).

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.